

Improving Health Care Response to Preeclampsia

Maurice L. Druzin^a, MD, Laurence E. Shields,^b MD, Nancy L. Peterson,^{ac} RNC, PNNP, MSN, Kathryn Melsop^{ac}, MS, Valerie Cape^{ac} (Eds).

Stanford University School of Medicine^a; Dignity Health^b; California Maternal Quality Care Collaborative^c

Suggested citation:

Maurice L. Druzin, MD, Laurence E. Shields, MD, Nancy L. Peterson, RNC, PNNP, MSN, Kathryn Melsop, MS, Valerie Cape, BSBA. Preeclampsia Toolkit: Improving Health Care Response to Preeclampsia (California Maternal Quality Care Collaborative Toolkit to Transform Maternity Care) Developed under contract ##08-85012 with the California Department of Public Health; Maternal, Child and Adolescent Health Division; Published by the California Maternal Quality Care Collaborative, August 2013.

Funding for the development of this toolkit was provided by:

Federal Title V MCH block grant funding from the California Department of Public Health; Maternal, Child and Adolescent Health Division and Stanford University.

The California Toolkit to Transform Maternity Care called “Improving the Health Care Response to Preeclampsia” was reviewed by the California Department of Public Health; Maternal Child and Adolescent Health Division. This toolkit is considered a resource, but does not define the standard of care in California. Readers are advised to adapt the guidelines and resources based on their local facility’s level of care and patient populations served and are also advised to not rely solely on the guidelines presented here.

Copyright information

© 2013 California Department of Public Health. The material in this toolkit may be reproduced and disseminated in any media in its original format, without modification, for informational, educational and non-commercial purposes only. A nominal sum to cover costs of reproduction and distribution can be assessed. Any modification or use of the materials in any derivative work is prohibited without prior permission of the California Department of Public Health.

For correspondence, please contact:

CMQCC

Nancy Peterson, RNC, PNNP, MSN, IBCLC
Managing Editor, Transforming Maternity Care Series
1265 Welch Road
Stanford, CA 94305
Phone: (650) 723-4849
Fax: (650) 721-5751
Email: nursenp@stanford.edu
Website: <http://www.cmqcc.org/>

CA Department of Public Health

Maternal, Child and Adolescent Health Division

Leona Shields, PHN, MN, NP
1615 Capitol Ave., MS 8306
PO Box 997420
Sacramento, CA 95899-7420
Phone: (916) 650-0368
Fax: (916) 650-0304
Email: Leona.Shields@cdph.ca.gov
Website: <http://www.cdph.ca.gov>

ACKNOWLEDGEMENTS

CMQCC would like to thank the California Department of Public Health; Maternal Child Health Division leaders Shabbir Ahmad, DVM, MS, PhD and Leona Shields, PHN, MN, NP, for their leadership to improve maternal health in California.

CMQCC would like to thank volunteer members of the Preeclampsia Task Force (PTF), the Task Force Co-Chairs, Maurice Druzin, MD, and Laurence Shields, MD, for their contributions to the Preeclampsia Care Guidelines and Compendium of Best Practices.

Preeclampsia Task Force

Co-Chairs:

- Maurice Druzin, MD – Stanford University School of Medicine
- Nancy Peterson, RNC, PNNP, MSN – Stanford University School of Medicine
- Laurence Shields, MD – Marian Regional Medical Center/Dignity Health

- Thomas Archer, MD – University of California, San Diego Medical Center
- Ocean Berg, RN, MSN, IBCLC – Nurse Family Partnership Program, MCH
- Claire Brindis, PhD – University of California, San Francisco
- Brenda Chagolla, RN, MSN, CNS – University of California Davis Medical Center
- Holly Champagne, MSN, RNC, CNS – Kaiser Permanente, Roseville
- Meredith Drews – Preeclampsia Foundation
- Racine Edwards-Silva, MD, FACOG – Olive View UCLA Medical Center
- Kristi Gabel, RNC-OB, C-EFM, MSN, CNS – Sutter Roseville Medical Center
- Dana Hughes, DrPh – University of California, San Francisco
- Thomas Kelly, MD, University of California, San Diego School of Medicine
- Sarah Kilpatrick MD, PhD – Cedars-Sinai Medical Center
- Richard H. Lee, MD – University of Southern California Keck School of Medicine
- Audrey Lyndon, PhD, RNC, FAAN - University of California, San Francisco
- Elliott Main, MD – California Pacific Medical Center/CMQCC Medical Director
- Mark Meyer, MD, FAAEM – Kaiser Permanente, San Diego
- Connie Mitchell, MD, MPH – Maternal, Child and Adolescent Health
- Leona Shields, PHN, RN, NP, MFT - Maternal, Child and Adolescent Health
- Eleni Tsigas – Preeclampsia Foundation
- Linda V. Walsh, CNM, MPH, PhD, FACNM
- Mark Zakowski, MD – Cedars-Sinai Medical Center

Conflict of Interest: The contributing authors and reviewers do not have any affiliations or financial involvement that conflict with the material or recommendations presented in this toolkit.

CMQCC EXECUTIVE COMMITTEE

- Berneva Adams, MD – Southern California Kaiser Permanente Medical Group
- Shabbir Ahmad, DVM, MS, PhD – MCAH, California Department of Public Health
- Lisa Bollman, RNC, MSN, CPHQ - Community Perinatal Network
- James Byrne, MD – Santa Clara Valley Medical Center
- Leslie Cragin, CNM, PhD, FACNM
- Arlene Cullum, MPH – Sutter Medical Center, Sacramento
- Tracy Flanagan, MD – Kaiser Permanente Medical Group, Richmond
- Bill Gilbert, MD – Sutter Health, Sacramento Sierra Region
- Jeffrey Gould, MD, MPH – CPQCC, Stanford University
- Tipu Khan, MD – University of Southern California, CAFFP
- Lisa Korst, MD – Childbirth Research Associates, LLC
- Leslie Kowalewski – March of Dimes
- David Lagrew, MD – Saddleback Memorial Medical Center
- Audrey Lyndon, PhD, RNC, FAAN – University of California, San Francisco
- Joe Parker, PhD – Office of Statewide Planning and Development Healthcare Outcomes
- Steven Parry, MD, FACOG, MCBWARD, MediCal Benefits Branch
- Manuel Porto, MD, University of California, Irvine
- Diana Ramos, MD, MPH – County of Los Angeles Public Health
- Usha Ranji, MS – Kaiser Family Foundation
- Leona Shields, PHN, RN, NP, MFT – MCAH, California Department of Public Health
- Stephanie Teleki, MPH, PhD - California HealthCare Foundation
- John Wachtel, MD – Stanford University School of Medicine
- Randy Winter, MD – Doctors Medical Center of Modesto, ACOG
- Teresa Wray, MD Kaiser Permanente Southern California

CMQCC

- Elliott Main, MD – Medical Director
- Anisha Abreo, MPH – Data Analyst
- Valerie Cape - Program Administrator
- Kathryn Melsop, MS – Program Manager
- Christine Morton, PhD – Program Manager
- Nancy Peterson, RNC, PNNP, MSN – Clinical Program Manager
- Barb Murphy, MS, RN – Director, Perinatal Programs

EXECUTIVE SUMMARY

Between 1999 and 2008, the rate of maternal deaths in California nearly doubled from 8 to 14 per 100,000 live births. In African-American women the rate was approximately four times higher than in all other racial/ethnic groups.¹

Hypertensive disorders of pregnancy are a leading cause of maternal mortality occurring in 12-22% of pregnancies.¹ These disorders are responsible for approximately 17% of maternal mortality in the United States.²⁻⁴ The California Pregnancy Associated Mortality Review (CA-PAMR) from 2002-2004, found a similar incidence of maternal mortality related to preeclampsia and associated syndromes.¹ The overall mortality rate for preeclampsia among the CA-PAMR deaths is 1.6/100,000. In addition, these disorders are one of the leading contributors to premature birth leading to significant neonatal morbidity and mortality.¹ Tragically, all of these deaths were determined to have at least some chance to alter the outcome, and half of those were determined to have a strong-to-good chance to alter the outcome.

A major quality improvement theme that emerged in the analysis of CA-PAMR preeclampsia cases was that despite triggers that clearly indicated a serious deterioration in the patient's condition, health care providers failed to recognize and respond to these signs in a timely manner leading to delays in diagnosis and treatment. In addition, the Preeclampsia Task Force recognized the need to emphasize additional educational efforts on the importance of accurate blood pressure measurement and initiating antihypertensive medications early and aggressively to prevent progression of the disease.

In response to the alarming trends in maternal morbidity and mortality and inconsistent application of evidence based guidelines for managing hypertensive disorders, the California Maternal Quality Care Collaborative (CMQCC) and the Preeclampsia Task Force developed the toolkit, "Improving Health Care Response to Preeclampsia." The toolkit authors represent a multi-disciplinary team of experts from every corner of the state and from both large and low volume Obstetric (OB) units. The editorial process in developing the toolkit was extensive and included peer review and consensus among experts from around the state. One of the most important goals of this project was to provide tools and guidelines gleaned from the "lessons learned" from the CA-PAMR review to assist others in decreasing maternal morbidity and mortality in the future.

Many of the tools and best practices outlined in the toolkit have been developed and recommended by national and international organizations including: American Congress of Obstetricians and Gynecologists (ACOG), National Institute for Health and Clinical Excellence (NICE), United Kingdom, Society for Obstetricians and Gynaecologists of Canada (SOGC). An update from the ACOG Hypertension in Pregnancy Workgroup in

the form of a major revision to the technical bulletin is planned for release this calendar year.

The toolkit provides a series of articles on best practices for hypertensive disorders that range in topic from diagnostic challenges to appropriate implementation of accepted medical therapy and recognition of institutional limitations in providing care for these complex maternal patients. Of particular interest, the toolkit addresses the management of severe preeclampsia < 34 weeks, the importance of recognition and treatment of delayed postpartum preeclampsia/eclampsia in the emergency department and early postpartum follow-up upon discharge. In addition, the toolkit provides care guideline summaries (in checklist, flowchart and table chart formats); it is organized into the following sections:

- Compendium of Best Practices: eighteen articles on multiple topics around hypertensive disorders
- Appendices: Collection of all Care Guidelines including tables, charts and forms that are highlighted in Article Sample forms for policy and procedure
- Slide set for Professional Education: slides that summarize the problem of and the best practices for preeclampsia to be used for local education and training

CMQCC and the California Department of Public Health (CDPH), Maternal, Child and Adolescent Health (MCAH) Division collaborated to develop and disseminate this toolkit using Title V MCH funds provided by CDPH-MCAH. The goal of this toolkit is to guide and support obstetrical providers, clinical staff, hospitals and healthcare organizations to develop methods within their facilities for timely recognition and organized, swift response to preeclampsia and to implement successful quality improvement programs for preeclampsia that will decrease short- and long-term preeclampsia-related morbidity in women who give birth in California.

REFERENCES

1. *The California Pregnancy-Associated Mortality Review. Report from 2002 and 2003 Maternal Death Reviews: California Department of Public Health, Maternal Child and Adolescent Health Division.* Sacramento 2011.
2. American College of Obstetricians and Gynecologists. Diagnosis and management of preeclampsia and eclampsia; ACOG Practice Bulletin No. 33. *Obstet & Gynecol.* January 2002, Reaffirmed 2010;99:159-167.
3. Clark SL, Belfort MA, Dildy GA, Herbst MA, Meyers JA, Hankins GD. Maternal death in the 21st century: causes, prevention, and relationship to cesarean delivery. *Am J Obstet Gynecol.* Jul 2008;199(1):36 e31-35; discussion 91-32 e37-11.

-
4. Baker DW, Asch SM, Keesey JW, et al. Differences in education, knowledge, self-management activities, and health outcomes for patients with heart failure cared for under the chronic disease model: the improving chronic illness care evaluation. *J Card Fail.* Aug 2005;11(6):405-413.

Draft

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	2
EXECUTIVE SUMMARY	4
HOW TO USE THIS TOOLKIT	10
BACKGROUND, PREPARATION AND MANAGEMENT	10
INTRODUCTION AND HISTORICAL PERSPECTIVE.....	12
CLINICAL PEARLS	15
CLASSIFICATION AND DIAGNOSIS OF HYPERTENSIVE DISORDERS OF PREGNANCY	19
SUSPECTED PREECLAMPSIA ALGORITHM	24
PATIENT CARE AND TREATMENT RECOMMENDATIONS	25
ACCURATE BLOOD PRESSURE MEASUREMENT.....	25
EARLY RECOGNITION	28
PREECLAMPSIA EARLY RECOGNITION TOOL (PERT) 30	
PROTEINURIA.....	31
ANTE, INTRA, POSTPARTUM NURSING MANAGEMENT AND ASSESSMENT OF PREECLAMPSIA: MATERNAL/FETAL ASSESSMENT AND MONITORING RECOMMENDATIONS.....	35
OUTPATIENT MANAGEMENT OF PREECLAMPSIA.....	39
CHRONIC HYPERTENSION IN PREGNANCY	41
ANTIHYPERTENSIVE AGENTS IN PREECLAMPSIA	45
MAGNESIUM SULFATE.....	48
ECLAMPSIA ALGORITHM 55	
SAMPLE PREECLAMPSIA/ECLAMPSIA MEDICATION BOX 56	
TEAMWORK AND COMMUNICATION	57
THE ROLE OF MEDICAL SIMULATION.....	68
FLUID MANAGEMENT IN PREECLAMPSIA.....	70
AIRWAY MANAGEMENT IN PREGNANT OR POSTPARTUM WOMEN HAVING SEIZURES	73
SPECIAL CIRCUMSTANCES: SEVERE PREECLAMPSIA AT < 34 WEEKS	76
CONSULTATION TRIGGERS IN SEVERE PREECLAMPSIA FOR ALL OBSTETRIC UNITS.....	85

POSTERIOR REVERSIBLE ENCEPHALOPATHY SYNDROME (PRES)	87
EMERGENCY DEPARTMENT and NON-OBSTETRICAL VISITS	92
EMERGENCY DEPARTMENT RECOGNITION and TREATMENT: FOCUS ON DELAYED POSTPARTUM PREECLAMPSIA and ECLAMPSIA	92
Part 1 of 2: Diagnosis - Evaluation and Treatment of Antepartum and Postpartum Preeclampsia and Eclampsia in the Emergency Department.....	94
Part 2 of 2: Treatment - Evaluation and Treatment of Antepartum and Postpartum Preeclampsia and Eclampsia in the Emergency Department.....	95
EDUCATION AND PATIENT INFORMATION	96
PRENATAL AND POSTPARTUM PATIENT COUNSELING OR EDUCATION* ...	96
APPENDICES	102
Algorithms: Appendices A-F.....	102
Appendix A: Sample Management of Eclampsia Algorithm	102
Appendix B: Sample Treatment of Severe Preeclampsia Algorithm	103
Appendix C: Suspected Preeclampsia Algorithm	104
Appendix D: Preeclampsia Early Recognition Tool (PERT)	105
Appendix E: Eclampsia Algorithm	106
Appendix F: Evaluation and Treatment of Antepartum and Postpartum Preeclampsia and Eclampsia in the Emergency Department Part 1 AND 2	107
Simulations/Drills: Appendices G-N	109
Appendix G: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: General Information	109
Appendix H: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: Learning Objectives	110
Appendix I: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: Patient Background Information	111
Appendix J: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: Equipment/ Materials List	112
Appendix K: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: Program Algorithm & GUI Notes	113
Appendix L: Severe Preeclampsia/Eclampsia In LDR v2.0 SimMan 3G: Debriefing Objectives	114
Appendix M: Severe Preeclampsia/Eclampsia In LDR v2.0 SimMan 3G: Debriefing Guide/Evaluation	115
Appendix N: Simulation Scenario: Hypertension in Pregnancy, Help with Seizure	116
Patient Education: Appendices O-Q.....	120

Appendix O: Preeclampsia Foundation Signs and Symptoms Information Sheet	120
Appendix P: SAMPLE: Discharge Instructions Following Delivery with Diagnosed Preeclampsia	121
Appendix Q: SAMPLE: Discharge Sheet for Preeclampsia, Eclampsia and HELLP Syndrome Patients	122
Miscellaneous: Appendices R-W.....	123
Appendix R: Prenatal Care Monitoring: Documentation of Diagnosis and Accurate ICD-9 Coding	123
Appendix S: Sample Preeclampsia/Eclampsia Medication Toolbox List	126
Appendix T: Special Considerations: Amphetamine/Cocaine Drug abuse and hypotension	127
Appendix U: Sample Nursing Management Policy and Procedure	130
Appendix V: Daily Assessment for Delivery versus Continuing Pregnancy	135
Appendix W: Classification of Evidence Grading	136

HOW TO USE THIS TOOLKIT

COMPENDIUM OF BEST PRACTICES

The Compendium of Best Practices consists of eighteen (18) articles authored by the CMQCC Preeclampsia Task Force. The articles highlight the current practices and recommendations for optimal care for hypertensive disorders that range in topic from diagnostic challenges to appropriate implementation of accepted medical therapy to recognition of institutional limitations in providing care for these complex maternal patients.

Most articles provide topic-specific tools for use at your facility. For example, the article titled, “Early Recognition” provides a tool to help clinicians recognize and respond to the evolving progression of preeclampsia. Each article also provides topic-specific recommendations for optimal care, evidence grading for the literature reviewed and full references.

PREECLAMPSIA CARE GUIDELINES

The Preeclampsia Care Guidelines provide summary information to assist with the identification and response to Preeclampsia/Eclampsia. One of the most important goals of this project was to provide tools and guidelines gleaned from the “lessons learned” from the CA-PAMR review to help others avoid similar maternal mortalities in the future.

SLIDE SET FOR PROFESSIONAL EDUCATION

A comprehensive slide set will be included in this toolkit for use by clinicians, educators and hospital administration. The slide set will include background information about the problem of hypertensive disorders and outline the key elements of the toolkit to provide support and guide providers, clinical staff, hospitals and healthcare organizations to develop strategies to improve early recognition and response to preeclampsia.

BACKGROUND, PREPARATION AND MANAGEMENT

The Best Practice articles that follow are topic-specific and include background information, current literature review and recommendations for clinicians who are preparing to respond to preeclampsia. In each article, a “grade” is provided for the level of evidence found in the literature to support the topic recommendations made by the Task Force. Many articles include specific tools—topic specific forms, tables or appendices—that are included at the end of each article.

The Compendium of Best Practices provides a broad range of background information for preeclampsia and provides the current literature and expertise from which the summary care guidelines were developed.

Table 1: Evidence Categories

Type of Study or Evidence	
I	Evidence obtained from at least one properly designed randomized controlled trial
II-1	Evidence obtained from well-designed controlled trials without randomization
II-2	Evidence obtained from well-designed cohort or case-control analytic studies, preferably from more than one center or research group
II-3	Evidence obtained from multiple time series with or without intervention. Well done QI studies with statistical process control analyses (or the like) fall into this category. Dramatic results in uncontrolled experiments also could be regarded as this type of evidence
III	Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees

Level of Recommendations	
A	Recommendations based on high quality and consistent evidence
B	Recommendations based on limited or inconsistent evidence
C	Recommendations based primarily on consensus and expert opinion

Adapted from United States Preventive Services Task Force (USPST) and ACOG

INTRODUCTION AND HISTORICAL PERSPECTIVE

Maurice Druzin, MD, Stanford University

Linda Walsh, CNM, MPH, PhD, FACNM

Laurence E. Shields, MD, Marian Regional Medical Center/Dignity Health

Nancy Peterson, RNC, MSN, PNNP, Stanford University

Hypertensive disorders of pregnancy are one of the most common medical disorders of pregnancy, occurring in 12-22% of pregnancies.¹ These disorders are responsible for approximately 17% of maternal mortality in the United States.²⁻⁴ The California Pregnancy Associated Mortality Review (CA-PAMR)³, found a similar incidence of maternal mortality related to preeclampsia and associated syndromes, such as severe preeclampsia, eclampsia and HELLP (Hemolysis, Elevated Liver enzymes, Low Platelet Count) from 2002-2004.. These disorders are responsible for approximately 17% of maternal mortality in the United States.²⁻⁴ The California Pregnancy Associated Mortality Review (CA-PAMR)¹, found a similar incidence of maternal mortality related to preeclampsia and associated syndromes from 2002-2004. In addition, hypertensive disorders are one of the leading contributors to premature birth leading to significant neonatal morbidity and mortality. Delivery of the fetus and placenta is an essential and critical component of the management of this disease. Induced premature delivery, placing the preterm newborn at significant risk, is often necessary to preserve the pregnant patient's health and life. The cost of preserving maternal health is a potential increase in the incidence of preterm delivery.

The association between preeclampsia and maternal mortality has been noted in the medical literature for over 150 years. Physicians in Great Britain and France identified the presence of albumin in the urine of pregnant women and edema as factors in the development of eclampsia in the mid-1800s. With the development of the sphygmomanometer in 1888, physicians recognized that elevated blood pressure was associated with eclampsia, leading to the triad of proteinuria, edema and hypertension being universally recognized as precursors to eclampsia. The term preeclampsia was introduced as a means to recognize the time period where delivery could potentially be used to prevent the progression to eclampsia. Organizational efforts to introduce prenatal care as an intervention to prevent and treat conditions of pregnancy that contributed to high maternal mortality rates largely addressed the identification and treatment of preeclampsia and eclampsia. The current pattern of prenatal care visits closely reflects the recommendations published by the Children's Bureau in 1924.⁴

Major emphasis has been placed on the prevention of eclamptic seizures, which are associated with a significant increase in both maternal and neonatal morbidity and mortality. The use of magnesium sulfate to prevent and treat seizures has been well accepted throughout the world as the standard of care.⁵⁻⁸ The current standard of

practice is to administer magnesium sulfate for seizure prophylaxis in patients with severe preeclampsia. Seizure prophylaxis for treatment of mild preeclampsia remains controversial.

Historically, less emphasis has been placed on the control of blood pressure to prevent stroke, and this aspect of management has recently been identified as a major gap in knowledge and application of proven therapeutic interventions. Treatment of systolic blood pressure at levels of equal to or greater than 160, and/or diastolic blood pressure of equal to or greater than 105 has been emphasized by Martin et al.⁹ However, the clinician may choose to institute therapy at lower levels of systolic and/or diastolic blood pressures. (Please refer to National Institute of Health and Clinical Excellence Guidelines).² The data from the CA-PAMR review from 2002-2004 confirmed the importance of this approach, as lack of timely therapeutic intervention at these levels of blood pressure was a consistent finding in patients dying of cerebrovascular accident (CVA) in the setting of preeclampsia/eclampsia.

There is little dispute that delivery of the fetus and placenta is the most important intervention in the treatment algorithm for preeclampsia/eclampsia. The aphorism that “delivery is the cure” is widely accepted in the obstetrical world, but it is clear that in many cases the multi-system pathology continues for a variable amount of time following delivery. There is greater recognition of the importance of continued evaluation in the postpartum period following delivery of the fetus and placenta, as serious clinical consequences persist for days and even weeks postpartum. These include severe levels of hypertension, onset of eclamptic seizures, and renal dysfunction. The onset of posterior reversible encephalopathy syndrome (PRES) is often diagnosed in the postpartum period. The initial two weeks postpartum seems to be a particularly vulnerable time for these complications, but reports of sequelae of preeclampsia have been reported up to six weeks postpartum.

The preeclampsia task force was charged with developing a toolkit to address the clinical spectrum of this serious disorder and develop systematic and evidence-based approaches to management of this disease. Maternal mortality is the unfortunate outcome of this disease in a considerable number of cases, but the morbidity and long-term effect on patients’ lives are significant, underreported, and underappreciated.^{10,11}

Data from the CA-PAMR and other state and international reports have emphasized the high rate of preventability of morbidity and mortality in 50-70% of cases of hypertensive disorders of pregnancy.^{4,10,12,13} The CA-PAMR analysis of maternal deaths from 2002-2004 revealed that all of maternal deaths related to preeclampsia had at least “some” chance of being altered by adherence to systematic, evidence-based and well-published algorithms for management of this disease. Almost half (48%) had a ‘good to strong’ chance to alter the outcome. The Quality Improvement Opportunities (QIOs) identified through CA-PAMR reinforced the international data suggesting increased preventability




Figure 1: Major Themes in QIOs among Preeclampsia Deaths, CA-PAMR 2002-2004

of mortality by following straightforward management approaches. The 3-Delay (3D) Model was developed in under-resourced countries to address the significant incidence of maternal mortality in those countries and is validated by current data. The “3 Delays” are: 1) delay in deciding to seek care, 2) delay in reaching care in time, and 3) delay in receiving adequate treatment.¹⁴ Unfortunately, these same delays are often present in our own state and country as reflected in Figure 1.

This toolkit has been developed to be applicable at all levels of hospitals that deliver care to pregnant women and newborn infants. The toolkit algorithms are straightforward and focus on critical action. They are flexible for use by a wide variety of practitioners who are likely to encounter women with hypertensive disorders of pregnancy including family practitioners, emergency department personnel, midwives, obstetricians, perinatologists, nursing staff, and labor and delivery staff. The focus of the toolkit is on the continuum of care from the prenatal period through delivery and postpartum.

CLINICAL PEARLS

Compiled by the Preeclampsia Task Force Committee

ACUTE TREATMENT:

- Antihypertensive medications administered within 1 hour and ideally within 30 minutes of arrival at a healthcare facility for blood pressures of 160 systolic, and/or 105-110 diastolic or greater is a critical initial step in decreasing morbidity and mortality.
- Magnesium sulfate therapy for seizure prophylaxis should be administered to any patients with:
 - Severe preeclampsia,
 - Preeclampsia with severe features, also known as atypical (subjective neurological symptoms such as headache or blurry vision or right upper quadrant or epigastric abdominal pain) AND
 - Should be considered in patients with mild preeclampsia (preeclampsia without severe features).
- Magnesium sulfate is the approved initial therapy for an eclamptic seizure.
- Algorithms for acute treatment of severe hypertension and eclampsia should be readily available or preferably posted in all labor and delivery units.
- Early post-discharge follow-up should be the norm for all patients diagnosed with preeclampsia/eclampsia. The Task Force recommends that follow-up occur within 3-7 days if blood pressure medication was used during the labor and delivery or postpartum and within 7-14 days if the diagnosis of preeclampsia was made but no medication was used.
- Postpartum patients presenting with hypertension, preeclampsia or eclampsia to the Emergency Department should be either assessed by or admitted to an obstetrical service. If they are treated in the Emergency Department and discharged, adequate follow-up must be arranged with an obstetrical provider.
- All institutions should consider preparing a severe preeclampsia/eclampsia box of medications and supplies needed for the treatment of preeclampsia (see Appendix) that includes at a minimum the following: Magnesium sulfate (including tubing, syringes and needles), Labetalol, Hydralazine and Calcium Gluconate. Additional medications such as second-line antihypertensives should be institution specific.

PATIENT ASSESSMENT:

- A high index of suspicion for hypertensive disorders of pregnancy and the syndrome of preeclampsia/eclampsia is required when encountering pregnant women with evidence of NEW ONSET hypertension and/or proteinuria.
- Preeclampsia is typically a disease of the late third trimester; however, earlier onset of preeclampsia prior to 34 weeks, is often more severe and may have an atypical presentation. This diagnosis should be considered in any patient with new onset symptoms and signs of hypertension and/or proteinuria.
- Patients presenting with vague symptoms such as headache, abdominal pain, shortness of breath, “I just don’t feel right,” or generalized swelling should be evaluated for atypical presentations of preeclampsia.
- Forty percent of patients with new onset hypertension or new onset proteinuria will develop preeclampsia.¹⁵
- Patients presenting with preeclampsia, severe preeclampsia or eclampsia to centers with limited resources to care for either the infant or mother should be stabilized and transferred to a center that has the capacity to care for expected complications of either the mother or infant.

PROVIDER and PATIENT EDUCATION:

- Healthcare professionals often tend to minimize signs and symptoms and therefore, may miss an opportunity to alter outcome.
- Use of patient education strategies, targeted to the educational level of the patients will increase patient awareness of signs and symptoms of preeclampsia.
- The importance of adequate prenatal care and access to obstetrical services should be emphasized for all socio-economic groups.
- Use of preeclampsia specific checklists, team training and implementation of a continuous process improvement strategy may reduce the morbidity associated with hypertensive disorders of pregnancy.
- The patient should be counseled that hypertensive disorders during pregnancy may predict future cardiovascular risk.
- There is no clinically validated screening strategy to predict the development of preeclampsia at this time.

REFERENCES

1. *The California Pregnancy-Associated Mortality Review. Report from 2002 and 2003 Maternal Death Reviews: California Department of Public Health, Maternal Child and Adolescent Health Division.* Sacramento 2011.
2. American College of Obstetricians and Gynecologists. Diagnosis and management of preeclampsia and eclampsia; ACOG Practice Bulletin No. 33. *Obstet & Gynecol.* January 2002, Reaffirmed 2010;99:159-167.
3. California Department of Health Care Services Maternal Child and Adolescent Health Branch. MCAH Bulletin California: The California Pregnancy-Associated Mortality Review (CA-PAMR) Report from 2002 to 2004 Maternal Death Reviews. 2012. <http://www.cdph.ca.gov/data/statistics/Documents/MO-CA-PAMR-MaternalDeathReview-2002-04.pdf>.
4. Thompson J, Walsh L, Merkatz I. The history of prenatal care: cultural, social and medical contexts. *New Perspectives on Prenatal Care.* 2008(1990):9-30.
5. Clark SL, Belfort MA, Dildy GA, Herbst MA, Meyers JA, Hankins GD. Maternal death in the 21st century: causes, prevention, and relationship to cesarean delivery. *Am J Obstet Gynecol.* Jul 2008;199(1):36 e31-35; discussion 91-32 e37-11.
6. Sibai BM. Magnesium sulfate prophylaxis in preeclampsia: Lessons learned from recent trials. *American Journal of Obstetrics and Gynecology.* Jun 2004;190(6):1520-1526.
7. Magpie Trial Collaborative Group. Do women with pre-eclampsia, and their babies, benefit from magnesium sulphate? *Lancet.* 2002;359:1877-1890.
8. Duley L, Galmezoghu A, Henderson-Smart D. Magnesium sulfate and other anticonvulsants for women with preeclampsia. *Cochrane Database of Systematic Reviews.* Vol 2003:2.
9. Martin J, Thigpen B, Moore R, Rose C, Cushman J, May W. Stroke and severe preeclampsia and eclampsia: a paradigm shift focusing on systolic blood pressure. *Obstet & Gynecol.* 2005;105(2):246-254.
10. World Health Organization. WHO Recommendations for Prevention and Treatment of Preeclampsia and Eclampsia: Evidence Base. 2011.
11. Aukes A, Wessel I, Dubois A, Aarnoudse J, Zeeman G. Self-reported cognitive functioning in formerly eclamptic women. *Am J Obstet Gynecol.* 2007;197:365.e361--365.e366.
12. CMACE. Saving Mothers Lives: reviewing maternal deaths to make motherhood safer: 2006-08. The Eighth Report on Confidential Enquiries into Maternal Deaths in the United Kingdom. *Br J Obstet Gynaecol.* 2011;118(Supplement 1):1-203.
13. Berg CJ, Callaghan WM, Syverson C, Henderson Z. Pregnancy-related mortality in the United States, 1998 to 2005. *Obstetrics and Gynecology.* Dec 2010;116(6):1302-1309.

-
14. Maternal United Nations Population Fund (UNFPA). Maternal Mortality Update 2002: A Focus on Emergency Obstetric Care. *New York:UNFPA.2003:8.*
 15. Barton JR, Sibai BM. Prediction and prevention of recurrent preeclampsia. *Obstet & Gynecology.* 2008;112(2 PART 1):359-372.

Draft

CLASSIFICATION AND DIAGNOSIS OF HYPERTENSIVE DISORDERS OF PREGNANCY

Maurice Druzin, MD, Stanford University

The diagnosis and classification of hypertensive disorders of pregnancy is primarily based on ACOG Practice Bulletin No. 33, January 2002.¹ The definition of hypertension in pregnancy is a blood pressure of ≥ 140 mm Hg systolic or ≥ 90 mm Hg diastolic. The criteria for the diagnosis of preeclampsia, which is a pregnancy specific syndrome usually occurring after 20 weeks of gestation, include new onset:

1. Blood pressure of 140 mmHg systolic or higher or 90 mmHg diastolic or higher that occurs after 20 weeks gestation in a women with previously normal blood pressure.
2. Proteinuria, defined as urinary excretion of 0.3 grams protein or higher in a 24-hour urine specimen.¹
3. Proteinuria, defined as urinary excretion of 0.3 grams protein or higher in a 24-hour urine specimen.¹

Preexisting hypertension prior to 20 weeks gestation would be considered chronic hypertension. Preexisting proteinuria prior to 20 weeks gestation would be suggestive of chronic renal disease.

New onset hypertension without proteinuria but with signs and symptoms of major end organ involvement such as headache, upper abdominal pain, hepatic dysfunction, pulmonary edema, or severe renal dysfunction, would potentially be indicative of “atypical preeclampsia.”²

The term gestational hypertension is used to describe cases in which elevated blood pressure without proteinuria develops in a woman after 20 weeks gestation and blood pressure levels return to normal postpartum (National High Blood Pressure working Group). As many as one quarter of women with gestational hypertension will develop proteinuria, i.e. preeclampsia.³

There is clearly potential for overlap of all these conditions, as a patient may present with gestational hypertension and progress to the preeclampsia/eclampsia syndrome very rapidly.

Table 1: Classification of hypertension in pregnancy

Chronic Hypertension	<ul style="list-style-type: none"> • BP of ≥ 140 mmHg systolic or 90 mmHg diastolic predating conception • Identified prior to 20 weeks gestation • Persists >12 weeks postpartum • Use of antihypertensive medications before pregnancy
Superimposed Preeclampsia or Eclampsia on Chronic Hypertension	<ul style="list-style-type: none"> • New onset proteinuria in a woman with hypertension prior to 20 weeks • Sudden increase in proteinuria if already present in early gestation • Sudden increase in BP • Development of HELLP syndrome • Development of headache, scotomata, or epigastric pain
Gestational Hypertension	<ul style="list-style-type: none"> • 140 mmHg systolic or ≥ 90 mmHg without proteinuria occurring after 20 weeks gestation • Transient diagnosis with normalization of BP by 12 weeks postpartum • May represent preproteinuric phase of preeclampsia or recurrence of chronic hypertension abated in mid-pregnancy • May evolve to preeclampsia • Retrospective diagnosis
Preeclampsia	<ul style="list-style-type: none"> • Occurring after 20 weeks of pregnancy • BP ≥ 140 mmHg systolic or ≥ 90 mmHg diastolic or higher • Proteinuria 0.3 grams protein or higher in a 24-hour urine specimen OR $\geq +1$ per dipstick OR P/C ratio > 0.3 mg/dL
Eclampsia	<ul style="list-style-type: none"> • Presence of new onset grand mal seizures in a pregnant woman with preeclampsia (rule out idiopathic seizure disorder or other central nervous system pathology such as intracranial hemorrhage, bleeding arteriovenous malformation, ruptured aneurysm) • New onset seizures 48-72 hours postpartum (other central nervous system pathology is the likely reason for the seizure after 7 days)
Severe Preeclampsia	<p>If one or more of the following criteria are present:</p> <ol style="list-style-type: none"> 1. Blood pressure of 160 mm Hg systolic or higher or 110 mm Hg diastolic or higher on two occasions at least 6 hours apart while the patient is on bed rest 2. Proteinuria of 5 grams or higher in a 24-hour urine specimen or 3+ greater on two random urine samples collected at least 4 hours apart 3. Oliguria of less than 500 mL in 24 hours 4. Cerebral or visual disturbances 5. Pulmonary edema or cyanosis 6. Epigastric or right upper-quadrant pain 7. Impaired liver function 8. Thrombocytopenia 9. Fetal growth restriction
HELLP Syndrome (subset of severe preeclampsia)	Hemolysis_Elevated Liver enzymes_Low Platelets

Adapted from ACOG Practice Bulletin #33, reaffirmed 2010 and from the National High Blood Pressure Education Program Working Group⁴

Table 2: Proposed ACOG HIP (Hypertension in Pregnancy) Working Group Recommendations from presentation at Society of Maternal Fetal Medicine (February 2013)

Current Terminology (ACOG Bulletin #33, 2002, reaffirmed 2010)	Proposed ACOG HIP Working Group Terminology (2013)
Mild preeclampsia (BP>140/90 mmHg)	<p>Preeclampsia without severe features</p> <ul style="list-style-type: none"> • The term “mild” will be eliminated • Presence of proteinuria is not required for the definition of preeclampsia IF there is hypertension AND any severe feature, such as sign(s) of end organ involvement which then qualifies for Severe Preeclampsia (see below)
Chronic Hypertension Gestational Hypertension Superimposed Preeclampsia	No Change in Definition
Severe preeclampsia (See Table 1)	<p>Preeclampsia with severe features (maternal end organ involvement) as listed in Table 1 except for the ELIMINATION of:</p> <ul style="list-style-type: none"> • Proteinuria of 5 g or greater in a 24 hour specimen • Fetal growth restriction

Other common associations with Preeclampsia:

HELLP Syndrome (Hemolysis, Elevated Liver enzymes, Low Platelets) is suggested when women with severe preeclampsia develop hepatic and hematologic manifestations as the predominant clinical picture, and is associated with an increased risk of adverse outcomes.^{3,5,6}

Chronic Hypertension complicating pregnancy is diagnosed by high blood pressure, BP \geq 140/90 or greater, known to predate conception. When preconception blood pressures are not known, elevated blood pressure detected before 20 weeks of gestation is often due to chronic hypertension. The most common etiology of chronic hypertension is most likely essential hypertension, although secondary hypertension as a result of renal disease, autoimmune disease, or vascular disease should be considered depending on the clinical presentation of the patient.

Superimposed Preeclampsia/Eclampsia chronic or gestational hypertension with superimposed preeclampsia is a common finding. Patients with underlying renal or

vascular disease have a high risk of developing superimposed preeclampsia, as do those with essential hypertension.

There have been a number of recommendations to further divide preeclampsia according to the gestational age of presentation into the following categories:

Less than 34 weeks gestation – early preeclampsia

Greater than 34 weeks gestation – late preeclampsia

Preeclampsia is either mild or severe under the currently accepted definitions. However, the ACOG HIP Task Force is recommending the elimination of the term “mild”. The recommendation is to use the terms “preeclampsia without severe features” or “preeclampsia with severe features.” The current rigid assignment of patients with this disease into a category of “mild preeclampsia” will often be detrimental to the appropriate management of patients. This disease is often not stable or static, but may evolve from “mild” preeclampsia to severe preeclampsia, HELLP Syndrome and/or eclampsia within a matter of hours. Rapid progression is typically seen in preeclampsia with onset prior to 34 weeks.⁷

EVIDENCE GRADING

Level of Evidence: II-2, II-3

REFERENCES


1. American College of Obstetricians and Gynecologists. Diagnosis and management of preeclampsia and eclampsia; ACOG Practice Bulletin No. 33. *Obstet & Gynecol.* January 2002, Reaffirmed 2010;99:159-167.
2. Sibai BM, Stella CL. Diagnosis and management of atypical preeclampsia-eclampsia. *American Journal of Obstetrics and Gynecology.* May 2009;200(5):481 e481-487.
3. Sibai B, Ramadan M, Chari R, Friedman S. Pregnancies complicated by HELLP syndrome (hemolysis, elevated liver enzymes, and low platelets): subsequent pregnancy outcome and long-term prognosis. *American Journal of Obstetrics and Gynecology.* 1995;172:125-129.
4. National High Blood Pressure Education Program Working Group. *American Journal of Obstetrics and Gynecology.* 2000;183:S1-S22.
5. Sibai B, Ramadan M, Usta I, Salama M, Mercer B, Friedman S. Maternal morbidity and mortality in 442 pregnancies with hemolysis, elevated liver enzymes, and low platelets (HELLP Syndrome). *American Journal of Obstetrics and Gynecology.* 1993;169:1000-1006.

-
6. Barton J, Sibai B. Hepatic imaging in HELLP syndrome (hemolysis, elevated liver enzymes, and low platelet count). *American Journal of Obstetrics and Gynecology*. 1996;174(1820-1825 discussion 1825-1827).
 7. Sibai BM. Evaluation and management of severe preeclampsia before 34 weeks' gestation. *Am J Obstet Gynecol*. Sep 2011;205(3):191-198.

Draft

SUSPECTED PREECLAMPSIA ALGORITHM

Suspected Preeclampsia Flowchart
 Diagnosis and Management



NOTE: There are no longer real differences in management between Preeclampsia and Gestational HTN in BP Management and decision to deliver. TREAT BP ACCORDINGLY and DELIVER for abnormal labs or symptoms

PATIENT CARE AND TREATMENT RECOMMENDATIONS


ACCURATE BLOOD PRESSURE MEASUREMENT

Kristi Gabel, RNC-OB, C-EFM, MSN, CNS, Sutter Roseville Medical Center

BACKGROUND

The current method used most often in the hospital setting for accurate measurement of blood pressure is the oscillatory method, or automated blood pressure machine, which tends to underestimate both systolic and diastolic readings by as much as 10 mm Hg^{1,2}. In the clinic setting and physician offices, blood pressure measurement is often used with the aneroid (mechanical type with a dial) sphygmomanometer. Refer to Table 1 for steps in obtaining accurate blood pressure measurement and Figure 1 for recommended cuff sizes.

Table 1: Steps for Obtaining Accurate Blood Pressure Measurements³

<p>Step 1: Prepare equipment</p>	<ul style="list-style-type: none"> a. Mercury sphygmomanometer is gold standard, can use validated equivalent automated equipment b. Check cuff for any defaults c. Obtain correct size cuff: width of bladder 40% of circumference and encircle 80% of arm (See Figure 1)
<p>Step 2: Prepare the patient:</p> 	<ul style="list-style-type: none"> a. Use a sitting or semi-reclining position with back supported and arm at heart level b. Patient to sit quietly for 5 minutes prior to measurement c. Bare upper arm of any restrictive clothing d. Patients feet should be flat, not dangling from examination table or bed, and her legs uncrossed e. Assess any recent (within previous 30 minutes) consumption of caffeine or nicotine. If blood pressures are at the level that requires treatment, consumption of nicotine or caffeine should not lead to delays in instituting appropriate anti-hypertensive therapies
<p>Step 3: Take measurement</p>	<ul style="list-style-type: none"> a. Support patients arm at heart level, seated in semi-fowlers position b. For auscultatory measurement: use first audible sound (Kortokoff I) as systolic pressure and use disappearance of sound (Kortokoff V) as diastolic pressure c. Read to the nearest 2 mmHg d. Instruct the patient not to talk e. At least one additional readings should be taken within 15 minutes f. Use the highest reading g. If greater than or equal to 140/90, repeat within 15-30 minutes and if still elevated, further evaluation for preeclampsia is warranted. <p>Do not reposition patient to either side to obtain a lower BP. This will give you a false reading.</p>
<p>Step 4: Record Measurement</p>	<p>Document BP, patient position, and arm in which taken</p>

Adapted from Peters RM (2008) High blood pressure in pregnancy. Nursing for Women's Health, Oct/Nov, pgs 410-422. Photo courtesy of and printed with permission by Kristi Gabel, RNC-OB, C-EFM, MSN, CNS, Sutter Roseville Medical Center

Figure 1: Recommended cuff sizes

Arm Circumference (cm)	Cuff Size
22-26	“Small Adult”: 12x22 cm
27-34	“Adult”: 16x30 cm
35-44	“Large Adult”: 16x36 cm
45-52	“Adult Thigh”: 16x42 cm




Photo courtesy of and printed with permission by Kristi Gabel, RNC-OB, C-EFM, MSN, CNS, Sutter Roseville Medical Center

Accurate blood pressure measurements in obese women can be quite challenging and it is extremely important to use an appropriate sized cuff. In women with an upper-arm circumference of more than 34cm, large adult cuffs or thigh cuffs can be used to improve blood pressure accuracy. For upper-arm measurements greater than 50cm, the American Heart Association recommends using a cuff on the forearm and feeling for the appearance of the radial pulse at the wrist to estimate systolic blood pressure. However, the accuracy of forearm measurement is not reliable.⁴

EVIDENCE GRADING

Level of Evidence: II and III

REFERENCES

1. Natarajan P, Shennan A, Penny J, Halligan A, de Swiet M, Anthony J. Comparison of auscultatory and oscillometric automated blood pressure monitors in the setting of preeclampsia. *American Journal of Obstetrics and Gynecology*. 1999;181 (5 Pt 1):1203-1210.
2. Ogedegbe G, Pickering T. Principles and techniques of blood pressure measurement. *Cardiology Clinics*. 2010;28(4):571-586.
3. Peters R. High blood pressure in pregnancy. *Nursing for Womens Health*. 2008;12(5410-421; quiz 422).

4. Pickering T, Hall J, Appel L, et al. Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public education of the American Heart Association Council on high blood pressure research. *Hypertension*. 2005;45:142.

Draft

EARLY RECOGNITION

Nancy Peterson, RNC, MSN, PNNP, Stanford University
 Laurence E. Shields, MD, Marian Regional Medical Center/Dignity Health

BACKGROUND

Early recognition and treatment of worsening signs and symptoms of preeclampsia was identified as a critical factor in reducing maternal morbidity and mortality in the 2002-2004 data from the California Pregnancy-Associated Maternal Mortality Review.¹ Of the 24 deaths due to preeclampsia, 48% (n=12) had a good-to-strong chance to alter the outcome, and an additional 48% (n=12) had some chance to alter the outcome. A major theme that emerged from the quality improvement data analysis was that despite clear triggers indicating a serious deterioration in the patient's condition, health care providers failed to recognize and respond to these clinical signs in a timely manner. In fact, missed vital sign "triggers" occurred in 60% of the preeclampsia deaths. Many of these cases documented elevated blood pressures in the medical record that should have alerted the health care team to implement an immediate clinical action to treat the women with antihypertensive medication. In addition, there were other "triggers" such as: proteinuria, headache, epigastric pain, deteriorating fetal status and altered mental status that were not recognized as serious or, in some cases, were dismissed or misinterpreted. In addition, the presence of various co-morbidities (pulmonary edema, hypotension, liver hematoma) was not considered in the context of the overall clinical picture. These cases demonstrated an overall lack of critical thinking or "putting the pieces of the puzzle" together to form a diagnosis by the health care team.

Another factor that can cause delays in early recognition, diagnosis and management of preeclampsia are the physiological changes of pregnancy, such as vasodilatation as a result of increased progesterone, that leads to a period of relative hypotension in the second trimester. Symptoms, such as headache, abdominal discomfort and "swelling" secondary to edema are common in normal pregnancy. Patients presenting with these symptoms may be judged to have "normal" changes of pregnancy and the signs of deterioration may therefore be masked for several hours. This can result in missed opportunities for timely interventions that may decrease the risk of adverse outcomes and/or death. Similar findings noted in the critical care literature, found that as many as 80% of patients have physiological abnormalities outside of the normal range in the 24 hours preceding critical care admission and up to 75% of such patients have at least one potentially life-threatening factor in the 8 hours before intensive care admission.²

The National Health System of the United Kingdom has been using a "Modified Obstetric Early Warning System" (MEOWS) chart ever since the 2003-2005 Confidential Enquiry into Maternal and Child Health (CEMACH) Report recommended its use.³ This MEOWS chart is designed to aid in the early recognition of severity of illness to allow prompt treatment of patients who are at risk for life-threatening complications.

In 2010, the Joint Commission also issued a Sentinel Alert, “Preventing Maternal Death”, recommending that all centers have a process in place for recognition and response to a patient’s deteriorating condition with written criteria describing early warning signs and indicating when to seek further assistance.⁴

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. All centers should develop:
 - a. a process for both the recognition and appropriate response in the event of a patient’s deteriorating condition.
 - b. Written criteria describing early warning signs and intervention strategies. Whenever possible, these should be built into the electronic medical record system. [See example of a Preeclampsia Early Recognition Tool (PERT) attached.]
 - c. Protocols and drills for recognizing, responding to and treating preeclampsia.

REFERENCES


1. *The California Pregnancy-Associated Mortality Review. Report from 2002 and 2003 Maternal Death Reviews: California Department of Public Health, Maternal Child and Adolescent Health Division.* Sacramento 2011.
2. Tarassenko L, Hann A, Yang D. Integrated monitoring and analysis for early warning of patient deterioration. *Br J Anesth.* 2006;97:64-68.
3. CMACE. Saving Mothers Lives: reviewing maternal deaths to make motherhood safer: 2006-08. The Eighth Report on Confidential Enquiries into Maternal Deaths in the United Kingdom. *Br J Obstet Gynaecol.* 2011;118(Supplement 1):1-203.
4. The Joint Commission. Preventing Maternal Death. Sentinel Event Alert. Issue 44. 2010;
http://www.jointcommission.org/sentinel_event_alert_issue_44_preventing_maternal_death. Accessed January 26, 2010.

PREECLAMPSIA EARLY RECOGNITION TOOL (PERT)

Preeclampsia Early Recognition Tool (PERT)

ASSESS	NORMAL (GREEN)	WORRISOME (YELLOW)	SEVERE (RED)
Awareness	Alert/Oriented	• Agitated/Confused • Drowsy • Difficulty speaking	• Unresponsive
Headache	None	• Mild headache • Nausea, Vomiting	• Unrelieved Headache
Vision	None	• Blurred or impaired	• Temporary blindness
Systolic BP (mmHG)	100-139	140-159	≥160
Diastolic BP (mmHG)	50-89	90-105	≥105
HR	61-110	111-129	≥130
Respiration	11-24	25-30	<10 or >30
SOB	Absent	Present	Present
O2 Sat (%)	≥95	91-94	≤90
Pain: Abdomen or Chest	None	• Nausea, Vomiting • Chest Pain • Abdominal Pain	• Nausea, Vomiting • Chest Pain • Abdominal Pain
Fetal Signs	• Category I • Reactive NST	• Category II • IUGR • Non-Reactive NST	• Category III
Urine Output (ml/hr)	≥50	30-49	≤30 (in 2 hrs)
Proteinuria	Trace	• +1/+2 • ≥300mg/24 hours	• >+3 • ≥5g/24hrs*
Platelets	>100	50-100	<50
AST/ALT	<70	>70	>70
Creatinine	<0.8	0.9-1.2	>1.2
Magnesium/Mag Sulfate Toxicity	• DTR +1 • Respiration 16-20	• Depression of patellar reflexes	• Respiration <12

*Level of Proteinuria is not an accurate predictor of pregnancy outcome



6.19.13.v1

Adapted from the Modified Obstetric Early Warning System (MEOWS) in "Saving Mothers Lives: reviewing maternal deaths to make motherhood safer (2003-2005). The Seventh Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom 2007

PROTEINURIA

Holly Champagne, MSN, RNC-OB, CNS, Kaiser Roseville Permanente Medical Center
Sarah Kilpatrick, MD, PhD, Cedars Sinai Medical Center

BACKGROUND

Proteinuria may be identified and quantified by chemstick “dip”, timed urine collection or protein/creatinine ratio. The amount of protein present is one determinant of severe preeclampsia (Table 1). A 24-hour urine collection is considered the gold standard for quantification of urine protein.^{1,2} Past studies identified inconsistent results from one-time urine chemstick tests.^{3,4} However, recent articles confirmed the accuracy and value of urine dipstick tests as a method to assess proteinuria.^{5,6} Timed urine collections of 2-hours (h), 4-h, and 12-h have shown to correlate well with 24-h urine results and may be an acceptable alternative when there is not enough time to collect a 24-h urine.⁷⁻⁹ Chemstick testing of urine, while approved as a FDA Clinical Laboratory Improvement Amendments (CLIA) waived test, nevertheless requires considerable resources to meet the College of American Pathologists (CAP) accreditation standards.¹⁰ Some facilities routinely screen all patients who arrive for evaluation in triage with a chemstick dip while others send a sample for urinalysis to the laboratory for evaluation. Either method of assessment appears to be adequate for initial patient assessment.

The urine protein/creatinine ratio is a measurement designed to compensate for the variation in protein concentration in urine by comparing the amount of protein to the concentration of creatinine present (Figures 1, 2). Recent recommendations do not prove one method of urine testing for proteinuria superior to another and each is acceptable to use. Clinicians need to be aware of what testing methods are used by the laboratories in their practice setting(s).

Similar to gestational hypertension, isolated new onset proteinuria, in the absence of urinary tract infection (UTI), is associated with a significantly increased risk for development of preeclampsia/atypical preeclampsia.¹¹

Table 1. Proteinuria Values in Mild and Severe Preeclampsia

Preeclampsia	Dipstick	24-hour urine (UNIT)	Protein: Creatinine ratio (See Figures 2,3 for how to calculate)
Mild	1-2+	≥ 0.3 gm/24 hours	≥ 0.3
Severe	≥3+	≥5 grams/day	Not defined

For diagnosis: Urine dipstick samples obtained twice, four hours apart and in absence of infection.^{1,2}

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. Urine dipstick is an acceptable initial screen. If positive (1+ or more) further evaluation is warranted. The 24-hour urine collection is the gold standard and should be used to confirm significant proteinuria if time allows.
2. Alternative testing may include 2-, 4-, 12-hour urine collection or protein/creatinine ratio, if these methods are available for use at the institution.
3. The amount of proteinuria should be used to classify preeclampsia as mild or severe, and should not be otherwise used to predict severity of disease or guide patient management.^{12,13} (Refer to Special Circumstances: Severe Preeclampsia at <34 weeks for additional considerations for <34 week gestation).
4. The level of proteinuria should not be used as the only criterion for management decisions.
5. Do not delay care in order to obtain 24-hour urine results; proceed with treatment/management if other criteria for severe preeclampsia are present.
6. Obtain baseline 24-hour urine protein or validated equivalent from those patients with proteinuria present in early or pre-pregnancy. Use heightened surveillance, carefully evaluate for symptoms of severe preeclampsia, and monitor for an increase in excreted protein in this population.
7. Preeclampsia, eclampsia, severe gestational hypertension, and/or HELLP syndrome, may occur without proteinuria.^{10,12-15} Close monitoring remains necessary for non-proteinuric pregnant patients with blood pressures ≥ 140 systolic and/or ≥ 90 diastolic.
8. The presence of new onset proteinuria in the absence of elevated blood pressure requires careful and more frequent patient surveillance (weekly to twice weekly) for the possible development of preeclampsia.^{11,16,17}
9. The presence of new onset proteinuria in the absence of elevated blood pressure requires careful and more frequent patient surveillance (weekly to twice weekly) for the possible development of preeclampsia.^{11,16,17}

Figure 1: Calculation of creatinine ratio

<p>Calculation of urine protein: creatinine ratio</p> <p>$(Urine\ protein \times 0.088) \div (urine\ creatinine)$</p>

Figure 2: Online calculator for creatinine ratio

<p>Online Calculator for urine protein: creatinine ratio</p> <p>http://www.easycalculation.com/medical/urinaryprotein.php</p>
--

EVIDENCE GRADING

Level of Evidence: III-C

REFERENCES

1. American College of Obstetricians and Gynecologists. Diagnosis and management of preeclampsia and eclampsia; ACOG Practice Bulletin No. 33. *Obstet & Gynecol.* January 2002, Reaffirmed 2010;99:159-167.
2. Gifford R, August P, Cunningham G, Green L, et al. Working Group Report on High Blood Pressure in Pregnancy, No. 00-3029. *National Institutes of Health.* 2000(1-52).
3. Gangaram R, Ojwang P, Moodley J, Maharai D. The accuracy of urine dipsticks as a screening test for proteinuria in hypertensive disorders of pregnancy. *Hypertension in Pregnancy* 2005;24:117-123.
4. Waugh J, Clark T, Dibakaran T, Khan K, Kilby M. Accuracy of urinalysis dipstick techniques in predicting significant proteinuria in pregnancy. *Obstetrics and Gynecology.* 2004;103:769-777.
5. Payne B, Magee L, Cote A, et al. PIERS proteinuria: Relationship with adverse maternal and perinatal outcome. *Journal Obstetrics and Gynaecology Canada.* 2011; 33:588-597.
6. Von Dadelszen P, Payne B, Li J, et al. PIERS Study Group. Prediction of adverse maternal outcomes in pre-eclampsia. *Lancet.* 2011;277:219-227.
7. Abebe J, Eigbefoh J, Okogbenin S, Eifediyi R, Okusanya B. Accuracy of urine dipsticks, 2-h and 12-h urine collections for protein measurement as compared with the 24-h collection. *Journal Obstetrics and Gynaecology.* 2008(28):496-500.
8. Soni S, Aggarwal N, Dhaliwal L, Wangkheimayum S, Menon M. Absent Circadian Rhythm of Proteinuria in Hospitalized Patients with Preeclampsia. *Hypertension in Pregnancy.* 2010;Sept 22:1-7.
9. Airoidi J, Weinstein L. Clinical significance of proteinuria in pregnancy. *Obstet Gynecol.* 2007;Survey 62:11-24.
10. College of American Pathologists. Big tests, small problems-POC documentation. 2009; http://www.cap.org/apps/cap.portal?_nfpb=true&.
11. Sibai BM, Stella CL. Diagnosis and management of atypical preeclampsia-eclampsia. *American Journal of Obstetrics and Gynecology.* May 2009;200(5):481 e481-487.
12. Sibai B. Diagnosis, controversies and management of HELLP syndrome. *Obstetrics & Gynecology.* 2004;103:981-991.
13. Sibai B. Maternal-perinatal outcome in 254 consecutive cases. *American Journal of Obstetrics and Gynecology.* 1990;163:1049-1054.
14. Thornton C, Makris A, Ogle R, Tooher J, Hennessey A. Role of proteinuria in defying preeclampsia: clinical outcomes for mothers and babies. *Clin Exp Pharmacol Physiol.* 2010;37(4):466-470.

15. Homer C, Brown M, Mangos G, Davis G. Non-proteinuric preeclampsia: a novel risk indicator in women with gestational hypertension. *Journal of Hypertension*. 2008;26:295-302.
16. Morikawa M, Yamada T, Yamada T, et al. Pregnancy outcome of women who developed proteinuria in the absence of hypertension after mid-gestation. *Journal of Perinatal Medicine*. 2008;36:419-424.
17. Morikawa M, Yamada T, Minakami H. Outcome of pregnancy in patients with isolated proteinuria. *Curr Opin Obstet Gynecol*. 2009;21:491-496.

Draft

ANTE, INTRA, POSTPARTUM NURSING MANAGEMENT AND ASSESSMENT OF PREECLAMPSIA: MATERNAL/FETAL ASSESSMENT AND MONITORING RECOMMENDATIONS

Brenda Chagolla, RNC, MSN, CNS, University of California Davis Medical Center
Ocean Berg, RN, MSN, IBCLC, Nurse Family Partnership Program
Kristi Gabel, RNC-OB, C-EFM, MSN, CNS, Sutter Roseville Medical Center

BACKGROUND

Antepartum management as an outpatient can be considered for select women who have mild preeclampsia, who have access to follow-up appointments and can adhere to the treatment plan.¹ Management of mild preeclampsia at term, severe preeclampsia, or those whose conditions have worsened will require frequent monitoring of blood pressure, urinary output, cardiac and respiratory status, and central nervous system status. Because early recognition of changes in maternal-fetal status is imperative, women with preeclampsia should be cared for by a nurse who is experienced with care of high-risk patients and has the experience to recognize worsening signs of preeclampsia. Specific preventable errors in contributing to maternal deaths include failure to control blood pressure for hypertensive women, and failure to adequately diagnose and treat pulmonary edema in preeclampsia.²⁻⁴

Maintaining a quiet, calm atmosphere and controlling environmental stressors are important for the patient and the family. Frequent updates for the family on the condition of the mother help them to maintain a focus on the mother and infant rather than on the illness.⁵ Postpartum preeclampsia/eclampsia can develop four to six (4-6) weeks after birth among women who had no evidence of preeclampsia during their pregnancy or at the time of delivery.⁶ Women should be given specific instructions prior to discharge on signs and symptoms that warrant immediate follow up.

KEY LEARNING POINTS

1. Assess for signs and symptoms of worsening or severe preeclampsia and notify provider if any of these are present:
 - Increasing blood pressure
 - Headache
 - Altered level of consciousness – agitation, restless, lethargy, hallucinations, confusion
 - Visual disturbances – blurred vision, floaters, spots, blind spot
 - Upper abdominal pain
 - Urine output <30 ml/hr
 - Shortness of breath
 - Complaints of chest pain
 - SaO₂ < 95%

- Cough
 - Tachypnea > 26 breaths per minute
 - Tachycardia > 100 bpm
 - Adventitious breath sounds
 - Eclamptic seizure
 - Magnesium toxicity
2. Patient care assignments should take into account the level and expertise of the clinician or nurse assigned to care. Patients diagnosed with severe preeclampsia should be staffed with a 1:1 nurse to patient ratio, with the most experienced nurse available.
 4. Women with severe preeclampsia should receive care by a multi-disciplinary team. The team should consist of an obstetric provider credentialed to perform cesarean sections, nursing, anesthesia, NICU, laboratory, blood bank, social work, and other sub-specialties as needed.
 5. Utilize the following as parameters (Table 1) as recommended guidelines for the frequency of nursing assessment. The recommended assessment frequencies listed in Table 1 are guidelines and additional or more frequent assessments can be done as needed based on patient condition.⁷

Table 1. Nursing Assessment Frequency

A. Mild Preeclampsia

	Mild Preeclampsia		
	Antepartum*	Intrapartum*	Postpartum*
BP, Pulse, Respiration, SaO2	Every 4 hours	Every 60 min	Every 4 hours
Lung sounds	Every 4 hours	Every 4 hours	Every 4 hours
Deep consciousness	Every 8 hours	Every 8 hours	Every 8 hours
Edema			
Assessment for headache, visual disturbances, epigastric pain			
Fetal status and uterine activity	Every shift	Continuous	N/A
Temperature	Per facility protocol		
Intake and output	Every 1 hour with totals every 8 and 24 hours		

*This is the minimum frequency recommended for the patient NOT on magnesium sulfate.

B. Severe Preeclampsia Nursing Assessment Frequency

	Severe Preeclampsia Intrapartum and Postpartum for women on Magnesium Sulfate
BP, Pulse, Respiration, SaO2	Every 5 mins during loading dose and q30 mins during maintenance of magnesium sulfate infusion. Can change to every 60 mins if any one or more of the following criteria are met: <ul style="list-style-type: none"> • Mild preeclampsia • BP stable without increases for a minimum of 2 hours • No antihypertensives within last 6 hours • Antepartum patient • Latent phase of labor Continuous SaO2 during magnesium infusion for intrapartum. For postpartum patient, check with vital signs
Lung sounds	Every 2 hours
Deep tendon reflexes & clonus, Level of consciousness Edema Assessment for headache, visual disturbances, epigastric pain	Every 4 hours
Temperature	Per facility protocol
Intake and output	Intake: <ul style="list-style-type: none"> • IV solutions and medication drips should all be on a pump • Total hourly intake should be ≤ 125 ml/hr • NPO with ice chips or as permitted by practitioner Output: <ul style="list-style-type: none"> • Insert foley with urimeter Calculate hourly, end of shift, and 24-hour totals
Fetal status and uterine activity	Continuous fetal monitoring

C. Post Eclamptic Seizure and Magnesium Sulfate Toxicity

Post Eclamptic Seizure and Magnesium Sulfate Toxicity for Ante, Intra and Postpartum	
BP, Pulse, Respiration	Every 5 min until stable
O2 Sat & LOC	Every 15 min for a minimum of 1 hour
Fetal Assessment and Uterine Activity	Continuous

D. Acute BP Treatment with IV Medication

Acute BP Treatment with IV Medication: Ante, Intra and Postpartum	
BP, Pulse, Respiration	Every 5-15 min until stable
O2 Sat & LOC	Every 5-15 min for a minimum of 1 hour
Fetal Assessment and Uterine Activity	Continuous

EVIDENCE GRADING

Level of Evidence: III-C

REFERENCES

1. Sibai BM. Diagnosis and management of gestational hypertension and preeclampsia. *Obstetrics and Gynecology*. Jul 2003;102(1):181-192.
2. Turner J. Diagnosis and management of pre-eclampsia: an update. *International Journal of Women's Health*. 2010;2:327-337.
3. The Joint Commission. Preventing Maternal Death. Sentinel Event Alert. Issue 44. 2010;
http://www.jointcommission.org/sentinal_event_alert_issue_44_preventing_maternal_death. Accessed January 26, 2010.
4. Clark SL, Belfort MA, Dildy GA, Herbst MA, Meyers JA, Hankins GD. Maternal death in the 21st century: causes, prevention, and relationship to cesarean delivery. *Am J Obstet Gynecol*. Jul 2008;199(1):36 e31-35; discussion 91-32 e37-11.
5. Perry I, Beevers D. The definition of preeclampsia. *Br J Obstet Gynaecol*. 1994;101(7).
6. Yancey L, Withers E, Bakes K, Abbot J. Postpartum preeclampsia: emergency department presentation and management. *J Emerg Med*. 2011;40(4):380-384.
7. Eggleston N, Trojano N, Harvey C, Chez B. *Clinical care guidelines*. Philadelphia: Wolters Kluwer/Lippincott Williams & Wilkins; 2013.

OUTPATIENT MANAGEMENT OF PREECLAMPSIA

Sarah Kilpatrick, MD, PhD, Cedars Sinai Medical Center

BACKGROUND

Once a diagnosis of preeclampsia has been made based on new onset systolic blood pressure ≥ 140 mmHg and or diastolic blood pressure ≥ 90 mmHg, and new onset significant proteinuria, or signs and symptoms of preeclampsia as seen in the Chapter: Classification and Diagnosis of Hypertensive Disorders of Pregnancy, Table 1, pg 20, the provider must decide if the woman has mild preeclampsia or severe preeclampsia. Outpatient treatment should only be considered for women with mild preeclampsia at less than 37 weeks and only after confirming fetal wellbeing and maternal stability.¹ It is imperative in the initial evaluation to document the severity of preeclampsia and the following evaluation is recommended: blood pressure, proteinuria assessment, CBC (complete blood count) with platelet count, AST (Aspartate Aminotransferase), ALT (Alanine Aminotransferase), Cr (Creatinine) and LDH (Lactate dehydrogenase).² The symptoms that should be assessed and documented as present or absent include headache, abdominal pain, and significant visual disturbances.

Fetal assessment should include NST (Non-stress Test) or BPP (Biophysical Profile), which includes NST plus fetal movement, tone, breathing, and heartrate and amniotic fluid volume, and ultrasound assessment of fetal growth. The goal of outpatient management in women with mild preeclampsia is early identification of the development of severe preeclampsia so that the woman is hospitalized and delivered if necessary, before significant maternal or fetal morbidity ensues.

If any abnormalities in either maternal or fetal assessments are consistent with severe preeclampsia, further management should occur in the hospital (see Chapter: Special Circumstances: Severe Preeclampsia At <34 weeks, pg 76). If mild preeclampsia is documented and outpatient management is considered then there should be a clear documented follow-up plan that is understood by the patient. Heightened surveillance is recommended to diagnose signs of worsening disease, which would prompt hospitalization and/or delivery that would generally includes twice-weekly maternal and fetal assessment. Maternal blood pressure, urine protein assessment and a verbal review of signs and symptoms should be performed twice per week. The fetus should have an NST and AFI (Amniotic Fluid Index) or BPP twice per week during outpatient observation. Additional maternal laboratory tests should be done as indicated if there is a suspicion of worsening disease. Once the patient develops any sign of severe preeclampsia *she should be admitted to the hospital and her plan should change accordingly*. If the patient continues to have only mild preeclampsia but reaches 37 weeks, the plan of treatment should include delivery. If the patient is diagnosed with severe preeclampsia she should be admitted to the hospital and—if gestational age is 34 weeks or greater—delivered.³ If she is less than 34 weeks with severe preeclampsia she should be admitted and

managed at a tertiary care facility with close observation for worsening disease or complications that necessitate delivery.

EVIDENCE GRADING

Level of Evidence: C

REFERENCES

1. American College of Obstetricians and Gynecologists. Diagnosis and management of preeclampsia and eclampsia; ACOG Practice Bulletin No. 33. *Obstet & Gynecol.* January 2002, Reaffirmed 2010;99:159-167.
2. Sibai BM. Diagnosis and management of gestational hypertension and preeclampsia. *Obstetrics and Gynecology.* Jul 2003;102(1):181-192.
3. Sibai BM. Evaluation and management of severe preeclampsia before 34 weeks' gestation. *Am J Obstet Gynecol.* Sep 2011;205(3):191-198.

CHRONIC HYPERTENSION IN PREGNANCY

Laurence E. Shields MD, Marian Regional Medical Center/Dignity Health

BACKGROUND:

The prevalence of chronic hypertension in the child-bearing age population has been reported to vary from 0.6% to greater than 22% depending on the age, BMI, and ethnicity of the patient, with about 1-5% of pregnant women having chronic hypertension.^{1,2} The rate of chronic hypertension in the pregnant population has increased significantly since 1999.² This increased incidence of chronic hypertension has been attributed to increasing maternal age, along with increased rates of obesity and diabetes. Pregnancies complicated by chronic hypertension are at increased risk for a variety of maternal complications³ including superimposed preeclampsia^{4,5}, pulmonary edema, placental abruption, acute renal failure, cerebral vascular accidents, stroke, and maternal death.^{2,6} The rate of many complications associated with superimposed preeclampsia appear to be similar to those noted with severe preeclampsia.⁷

Among women who died from a pregnancy-related cause in California between 2002-2004, 17% (N=25/145) died from preeclampsia. Among this cohort of pregnancy-related deaths, the true burden of hypertension was significantly higher.⁸ Fifty-seven of 145, or 39% of the women that died during pregnancy were noted to have hypertension during the prenatal period, at labor and delivery, or postpartum. This is five-to-six times higher than in the general obstetrical population in California, suggesting hypertension is a significant comorbidity for pregnancy-related deaths. Patients with a history of chronic hypertension also represented a significant percentage of those patients whom died from cardiomyopathy.

Chronic hypertension during pregnancy is defined as blood pressure (mmHg) ≥ 140 systolic or ≥ 90 diastolic prior to the 20th week of pregnancy (Table 1).^{1,9} It is preferable to identify chronic hypertension prior to 12 weeks gestation, in part because the normal nadir of maternal blood pressure during pregnancy occurs at approximately 16-18 weeks. A patient presenting with second trimester blood pressures slightly below the 140/90 cut-off for the diagnosis of chronic hypertension may in fact have mild to moderate chronic hypertension. Most national committees divide the severity of hypertension into mild or severe categories. The majority (90%) of patients diagnosed with chronic hypertension will have essential hypertension. The remaining 10% will have a secondary cause of the hypertension and patients presenting with severe hypertension first diagnosed in early pregnancy should be evaluated for secondary causes (pheochromocytoma, primary aldosteronism, Cushing Syndrome, sleep apnea, methamphetamine use, renal artery stenosis).^{1,10} As a result, it is possible that a patient's chronic hypertension increases with the risk for the development of superimposed preeclampsia and the presence of proteinuria at initial evaluation also increases the risk of adverse pregnancy outcomes.⁴ The most severe adverse outcomes of pregnancy related to hypertension (i.e., stroke and

cerebral vascular accidents) are most closely associated with systolic hypertension above 155-160 mmHg.¹¹

The most effective therapeutic approach to women with chronic hypertension during pregnancy is controversial. Treatment trials have been limited in size and yielded mixed results. Many experts argue that treatment of hypertension outside of pregnancy is directed towards reducing the longer-term risk of cerebral vascular and cardiac events, and the duration of pregnancy is unlikely to influence these outcomes in patients with mild chronic hypertension. Well-controlled randomized trials are limited in assisting clinicians in appropriate choices of medical therapy. Two Cochrane reviews are available detailing the results of treatment of mild and moderate chronic hypertension during pregnancy.^{7,12} In these reports, the use of beta-blockers and methyldopa were both associated with reductions in progression to severe hypertension, and beta-blockers were also associated with reductions in proteinuric preeclampsia, eclampsia, and neonatal Respiratory Distress Syndrome (RDS). Other widely accepted anti-hypertensive agents in pregnancy include labetalol, hydralazine, and nifedipine; however, there are some concerns that nifedipine may be associated with a modest increased risk for the development of superimposed preeclampsia.^{1,7} Neither of these reviews address the issue of the level of blood pressure control, and concerns have been raised that aggressive treatment may decrease placental perfusion and negatively impact fetal growth.¹²

Recommendations related to treatment have been primarily in the form of expert opinion and consensus recommendations from groups like American Congress of Obstetrics and Gynecology (ACOG), Society of Obstetricians and Gynaecologists of Canada (SOGC), and the National Institute for Health and Clinical Excellence (NICE) (Table 2). Women with chronic hypertension who are treated with medication should be closely supervised for both maternal and fetal status by a physician experienced in treating and monitoring hypertension in pregnancy.

Table 1. Diagnostic Criteria for Patients with Chronic Hypertension in Pregnancy among U.S. and Canada Organizations

Organization		Mild (mm Hg)		Severe (mmHg)
ACOG*	Systolic	140-159		≥160
	Diastolic	90-109		≥110
SOGC**	Systolic	≥ 140		≥160
	Diastolic	≥ 90		≥110
NICE***		Mild	Moderate	Severe
	Systolic	140-149	150-159	≥160
	Diastolic	90-99	100-109	≥110

*American College of Obstetrics and Gynecology.

**Society of Obstetricians and Gynaecologists of Canada.

***National Institute for Health and Clinical Excellence (NICE)

Table 2. Treatment Recommendations for Blood Pressure (Systolic/Diastolic mmHg) Patients with Chronic Hypertension among U.S. and Canada Organizations

Organization	Recommendation	Goal	Co-Morbidities****
ACOG*	150-160/100-105	<150/100	
SOGC**	140-159/90-109	130-155/90-109	130-139/80-90 mmHg
NICE***	150/100	<150/100	<140/90

* American College of Obstetrics and Gynecology.

** Society of Obstetricians and Gynaecologists of Canada

*** National Institute for Health and Clinical Excellence (NICE).

****Comorbid conditions (i.e., Special circumstances) are defined as the presence of impaired renal function, pre-gestational diabetes, cardiovascular disease, SOGC 2008;30 (3) S29

KEY LEARNING POINTS

Women with chronic hypertension should receive more frequent prenatal assessments during the late second and early third trimester due to the increased rate of maternal and fetal complications. The frequency of assessment should be weekly for those with stable blood pressure control, and every 3-4 days for those that are requiring increasing dosages of blood pressure medication.

1. Women with chronic hypertension should receive antihypertensive treatment if their blood pressure is in the severe range; they should be considered for therapy if their blood pressure elevation is mild to moderate, particularly if comorbid conditions such as diabetes, collagen vascular disease, or chronic renal disease are present.
2. Women presenting for their first prenatal visit in the mid-second trimester with blood pressures that are not quite high enough for a diagnosis of chronic hypertension (e.g., 130-139/80-89 mmHg) may in fact have chronic hypertension and should be observed more frequently for blood pressure exacerbation.
3. Cardiomyopathy should be part of the differential diagnosis and assessment for women presenting with symptoms of shortness of breath and chronic hypertension, particular if they are in a high-risk category (preexisting diabetes, collagen vascular disease, obesity, advanced maternal age, African American ethnicity, or long standing chronic hypertension).
4. Evaluation of cardiac function using echocardiography and laboratory assessment Brain Natriuretic Peptides (BNP), a test for cardiovascular disease, should be considered in these patients.

EVIDENCE GRADING

Level of Evidence: I-A, III-C

REFERENCES

1. ACOG. Chronic Hypertension in Pregnancy, Practic Bulletin #125. *Obstet Gynecol.* 2012;119:396-407.
2. Bateman BT, Bansil P, Hernandez-Diaz S, Mhyre JM, WM C, Kuklina E. Prevalence, trends, and outcomes of chronic hypertension: a nationwide sample of delivery admissions. *Am J Obstet Gynecol.* 2012;206:134 e131-138.
3. Rey E, Couturier A. The prognosis of pregnancy in women with chronic hypertension. *Am J Obstet Gynecol.* 1994;171:410-416.
4. Sibai B, Lindheimer M, Hauth J, et al. Risk factors for preeclampsia, abruptio placentae, and adverse neonatal outcomes among women with chronic hypertension. National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. *N Engl J Med.* 1998;339:667-671.
5. Sibai B. Chronic hypertension in pregnancy. *Obstet & Gynecology.* 2002;100.
6. Kuklina E, Meikle, S., Jamieson, D., Whiteman, M., Barfield, W., Hillis, S., Posner, S. Severe Obstetric Morbidity in the US, 1998-2005. *Obstetrics and Gynecology.* 2009;113:293-299.
7. Abalos E, Duley L, Steyn D, Henderson-Smart D. Antihypertension drug therapy for mild to moderate hypertension during pregnancy (Review). *Cochrane Database System Re* 2010;10. 2010.
8. California Department of Health Care Services Maternal Child and Adolescent Health Branch. MCAH Bulletin California: The California Pregnancy-Associated Mortality Review (CA-PAMR) Report from 2002 to 2004 Maternal Death Reviews. 2012. <http://www.cdph.ca.gov/data/statistics/Documents/MO-CA-PAMR-MaternalDeathReview-2002-04.pdf>.
9. Barton JR, Sibai BM. Prediction and prevention of recurrent preeclampsia. *Obstet & Gynecology.* 2008;112(2 PART 1):359-372.
10. ACOG. Committee Opinion no. 514: Emergent therapy for acute-onset, severe hypertension with preeclampsia or eclampsia. *Obstet Gynecol.* 2011;118:1465-1468.
11. Sibai B. Diagnosis, prevention, and management of eclampsia. *Obstet & Gynecol.* 2005;105(2):402-410.
12. Magee LA, Duley L. Oral beta-blockers for mild to moderate hypertension during pregnancy (Review). *Cochrane Database of Systematic Reviews.* 2009;2009:1:CD002863.

ANTIHYPERTENSIVE AGENTS IN PREECLAMPSIA

Tom Archer, MD, MBA, University of California San Diego

BACKGROUND

Early treatment of hypertension has consistently been found to reduce the incidence of hypertensive crisis, and data from multiple case series revealed increased rates of heart failure, pulmonary edema, stroke and death when antihypertensive medications were not used in women with severe gestational hypertension or severe preeclampsia.¹ According to ACOG, a hypertensive emergency is an acute-onset, severe hypertension that is persistent for 15 minutes or more.²

Treatment should be initiated for blood pressures that are ≥ 160 mmHg systolic or 105-110 mmHg diastolic.³ It should be noted that others have suggested that treatment should be initiated at a lower threshold of 155/105 if the primary goal is to reduce maternal intracranial hemorrhage, which remains the leading cause of death from preeclampsia.⁴

The goal of blood pressure control is not to return it to “normal” but rather to lower it to a range of 140-160/90-100, a level at which the risk of intracranial hemorrhage is reduced. Lowering the blood pressure below this range may reduce placental perfusion and harm the fetus.^{4,5} Below are key guidelines for the administration of antihypertensive medications.

KEY LEARNING POINTS

1. Antihypertensive therapy is reserved for women with systolic blood pressure greater than 160 mmHg or diastolic blood pressure greater than 105-110 mmHg. Increasingly, risk of stroke is felt to be correlated with maximum systolic blood pressure.^{6,7}
2. Hydralazine and labetalol are the two “first line” agents used for hypertension in preeclampsia. Hydralazine is an arteriolar dilator that reduces blood pressure but may cause tachycardia. Possible side effects are headache, risk of delayed maternal hypotension, which can be associated with fetal bradycardia, and rarely, upper abdominal (e.g., “epigastric”) pain, which may be confused with worsening preeclampsia. Labetalol is a combined alpha and beta-blocking agent, which reduces blood pressure by dilating arterioles and decreasing heart rate. Labetalol should be administered intravenously for acute hypertensive emergencies.²
3. Oral nifedipine (calcium channel blocker), IV esmolol (beta blocker) and IV nicardipine (calcium channel blocker) are second line drugs. Esmolol, is a very short-acting agent, and can cause the baseline fetal heart rate to decrease, but often resolves rapidly when esmolol is stopped.

4. Sodium nitroprusside is a very potent vasodilator that acts immediately and is rarely used. It must be used by experienced providers accompanied by invasive (e.g., an arterial line) blood pressure monitoring.
5. Consideration for consultation with anesthesiologists who are accustomed to the titration of vasoactive medications should be considered early for patients with uncontrolled blood pressure.
6. Placement of an arterial line may be helpful in women whose blood pressure is particularly difficult to control. There may also be cases where repeated blood studies will be necessary, e.g., magnesium levels, platelet counts, etc., and where repeated venipunctures may be difficult.
7. Women with severe hypertension requiring antihypertensive medications need to be observed carefully for signs of pulmonary congestion such as agitation, low oxygen saturation, cough, or rales on lung exam suggesting pulmonary edema or heart failure. Careful monitoring should be implemented.
9. Proper lateral positioning should be employed for these patients, since aortocaval compression can exacerbate uteroplacental insufficiency due to preeclampsia itself.
10. One of the most common and severe complications is hypertensive encephalopathy or Posterior Reversible Encephalopathy Syndrome (PRES).

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. Timely initiation of medication for treating elevated blood pressure is critical. The goal is to begin therapy within one hour and ideally within 30 minutes of presentation with severe range blood pressures. Initial therapy should consist of Labetalol 20 mg or Hydralazine 5-10 mg IV over 2 minutes. Hydralazine begins to have an effect within 5-20 minutes with its maximum effect occurring at 20-30 minutes. Labetalol onset is within 1-2 minutes and has its maximum effect after 5-10 minutes.
2. Maximum cumulative IV administered doses should not exceed the following:
Hydralazine 25 mg; labetalol 220 mg in 24 hours.⁷
3. The goal for blood pressure control is 140-160/90-100. Do not try to lower the blood pressure to “normal.”
4. The anesthesiologist should be seen as a resource for invasive blood pressure monitoring (if required) and medication titration.
5. An arterial line can be useful for acquiring multiple blood samples as well as for arterial pressure measurement.

EVIDENCE GRADING

Level of Evidence: III-C

REFERENCES

1. Magee LA, Abalos E, von Dadelszen P, Sibai B, Easterling T, Walkinshaw S. How to manage hypertension in pregnancy effectively. *Br J Clin Pharmacol*. Sep 2011;72(3):394-401.
2. ACOG. Committee Opinion no. 514: Emergent therapy for acute-onset, severe hypertension with preeclampsia or eclampsia. *Obstet Gynecol*. 2011;118:1465-1468.
3. ACOG. Diagnosis and Management of Preeclampsia and Eclampsia #33. *American College of Obstetricians and Gynecologists Practice Bulletin Number 33*. 2002 (Reaffirmed 2008).
4. Martin J, Thigpen B, Moore R, Rose C, Cushman J, May W. Stroke and severe preeclampsia and eclampsia: a paradigm shift focusing on systolic blood pressure. *Obstet & Gynecol*. 2005;105(2):246-254.
5. Creasy R, Resnik R, Iams J, Lockwood C, Moore T. *Creasy & Resnik's Maternal-Fetal Medicine: Principles and Practice*. Vol Sixth Edition, Chapter 35, pg. 673. 2008.
6. Chestnut D, Polley L, Tsen L, Wong C. *Chestnut's Obstetric Anesthesia: Principles and Practice, 4th Edition, Chapter 45, pg 984-986*. 2009.
7. American College of Obstetricians and Gynecologists. Diagnosis and management of preeclampsia and eclampsia; ACOG Practice Bulletin No. 33. *Obstet & Gynecol*. January 2002, Reaffirmed 2010;99:159-167.

MAGNESIUM SULFATE

Ocean Berg, RN, MSN, IBCLC, Nurse Family Partnership Program
Richard H. Lee, MD, University of Southern California
Brenda Chagolla, RN, MSN, CNS, University of California Davis Medical Center

BACKGROUND

Magnesium sulfate is the primary medication used in the prevention and management of eclamptic seizures and exerts its effect by depressing the central nervous system.¹

In the setting of severe preeclampsia, magnesium sulfate has been shown to significantly reduce the rate of eclampsia compared to placebo, phenytoin, nimodipine, or diazepam. In the setting of eclampsia, magnesium sulfate has been demonstrated to be superior to diazepam, phenytoin, or lytic cocktails in reducing the risk of recurrent seizures as well as reducing the risk of maternal death.²⁻⁷ However, controversy surrounds the use of magnesium in the setting of mild preeclampsia with some studies finding no difference and others alluding to a potential benefit.^{8,9} The Magpie Trial demonstrated the numbers needed to treat to prevent one seizure was 63 in subjects with severe preeclampsia and 109 in those with mild preeclampsia.² A similar rate of eclampsia (1.1%) was observed when magnesium sulfate was not used in women with less severe disease.¹⁰

Currently, the use of magnesium sulfate in severe preeclampsia and eclampsia is supported by The American Congress of Obstetricians and Gynecologists (ACOG),¹¹ World Health Organization (WHO),¹² Society of Obstetricians and Gynaecologists of Canada (SOGC),¹³ and the National Institute for Health and Clinical Excellence (NICE).¹⁴ Of these, the SOGC and WHO provide statements that support the use of magnesium sulfate in the setting of mild preeclampsia.^{12,13}

Magnesium sulfate is listed by the Institute of Safe Medication Practices (ISMP) as a high-alert medication.¹⁵ IN 2004 published data about Medication Errors in Labor, Delivery, Recovery, and Postpartum (LDRP's), magnesium sulfate was identified as the second most common product to be mismanaged in the Labor, Delivery and Recovery (LDR) area.¹⁵

In 2010 the Association of Women's Health Obstetric and Neonatal Nurses (AWHONN) updated their nurse staffing guidelines.¹⁶ Specific guidelines are identified for magnesium sulfate throughout the inpatient stay as follows:

- High-Risk Antepartum Care: A woman who is receiving IV magnesium sulfate should have one (1) nurse in continuous bedside attendance for the first hour of administration. Women receiving IV magnesium sulfate who are not in labor require a minimum of one (1) nurse to two (20) women.

- During labor of women with medical or obstetric complications: Women in labor who are receiving magnesium sulfate should have one (1) nurse in continuous bedside assistance for the first hour of administration and one (1) nurse to one (1) woman thereafter.
- Mother-Baby Care: Nurses caring for women receiving magnesium sulfate during the postpartum period should not have more than 1 other mother-baby couplet.

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. Mark magnesium sulfate as a high-alert medication.
2. Administer loading dose from an admixture bag and not from the maintenance solution.
3. Clearly label IV solutions, tubings and connections.
4. Use luer-lock connectors at all points.
5. Require independent verification by two nurses of magnesium sulfate IV pump infusion settings.
6. Standardize the ordering, storage, preparation, and administration of magnesium sulfate.
7. Improve access to information concerning indications, risks and benefits, and treatment of toxicity.

In 2011 the Joint Commission recommended that magnesium sulfate be written out and not abbreviated as MgSO₄. This can be misinterpreted as MS or MSO₄, which are abbreviations for morphine sulfate.¹⁷

Table 1: Steps for Preparation, Storage, Ordering and Administration of Magnesium Sulfate

Step 1: Preparation	<ul style="list-style-type: none"> a. Purchase commercially prepared standard concentrations of magnesium sulfate. b. Use standard premixed magnesium sulfate infusions in a volume different than Oxytocin to prevent accidental loading dose. c. Pharmacy should prepare any non-commercially prepared solutions. d. Use a piggy-back bag for magnesium sulfate loading doses, do not use the main infusion bag
Step 2: Storage	<ul style="list-style-type: none"> a. Label bags with a High Alert Sticker or distinctive colored label. b. Loading dose bags should be stored in a separate pharmacy area to prevent mix-ups.
Step 3: Order/ Transcribe	<ul style="list-style-type: none"> a. Use preprinted orders and order sets with the entire medication spelled out when prescribing magnesium sulfate. b. Utilize a High Alert warning on the automated dispensing machine (ADM) when magnesium sulfate is withdrawn.
Step 4: Administration	<ul style="list-style-type: none"> a. Label tubing used for infusing magnesium sulfate appropriately. b. An infusion pump should always be used. A ‘smart’ infusion pump with patient safety software activated should be used when available. <p>NOTE: An independent double check should be performed by two nurses when magnesium sulfate is initiated, dose is change and at change of shift.</p>
Step 5: Discontinuation	<ul style="list-style-type: none"> a. Disconnect tubing from main line immediately when magnesium sulfate is discontinued.

Table 2: Protocol for Administration of Magnesium Sulfate

A. Magnesium Sulfate Loading and Maintenance Dosage

	Loading (gm)	Infusion Rate (min)	Maintenance
Preeclampsia and Eclampsia	4-6	15-20	Infuse at 1-2 grams per hour via infusion pump
Recurrent Eclampsia	2	5	

B. Side Effects and Toxicity

Side Effects/Toxicity	Nursing Intervention
Cutaneous flushing, sweating, malaise, weakness, drowsiness	Keep room and patient cool (provide fan), educate patient about potential side effects; monitor patient movement and assist with getting out of bed
Transient decreased amplitude and frequency of contractions at the time of loading dose	Continuous external fetal and uterine monitoring*
Soreness at IV site	Warm soaks or ice site PRN
Decreased rate and depth of respiration, shortness of breath (SOB)	Discontinue treatment if SOB not relieved with oxygen
Diuresis	Strict Input & Output; document output per orders; magnesium sulfate is excreted exclusively in urine and an output of <30ml/hr may lead to magnesium toxicity
Disappearance of deep tendon reflexes	Notify physician if absent or significant change in baseline assessment
Heart block (decreased PR interval, increased QRS), chest pain	Avoid use in patients with cardiac conduction abnormalities
Pulmonary edema	Strict input and output, fluid restrict as ordered (usually 60-100ml/hr)

*Please see references cited.^{16,17}

Nursing care and assessment: (refer to Ante, Intra, Postpartum Nursing Management and Assessment of Preeclampsia: Maternal/Fetal Assessment and Monitoring Recommendations Section, pg. 35)

Increase frequency of assessments as indicated by patient condition.

If magnesium sulfate is unavailable, alternative anti-seizure medications such as a benzodiazepine (e.g., midazolam, lorazepam, or diazepam) or phenytoin should be used in the setting of eclampsia. Consultation with neurology is suggested to discuss continued medical prophylaxis for seizures if magnesium sulfate is unavailable. If a patient has mild preeclampsia and there is no magnesium sulfate available, we suggest observation alone.

KEY LEARNING POINTS

1. Magnesium sulfate dosage: Loading dose of 4-6 gm over 15-20 minutes. Maintenance dose of 1-2 g/hr.
2. Signs of magnesium toxicity: discontinue magnesium sulfate infusion and obtain a stat serum magnesium level in the following situations: respiratory rate <12

breaths/minute, complaint of shortness of breath, lethargy, nausea, vomiting, absent deep tendon reflexes, apnea, urine output <30cc/hour.¹

3. Magnesium levels: A specific therapeutic level is not known; however, serum magnesium levels may need to be monitored in certain circumstances (e.g., renal insufficiency, absent deep tendon reflexes). In such cases, magnesium levels between 4.8-8.4 mg/dL (4-7 mEq/L) is recommended. Symptoms of magnesium sulfate toxicity are seen with the following maternal serum concentrations: loss of deep tendon reflexes (9.6-12 mg/dL) (>7 mEq/L), respiratory depression (12-18 mg/dL) (>10 mEq/L), and cardiac arrest (24-30mg/dL) (>25mEq/L).
4. Calcium gluconate: the antidote for magnesium toxicity is calcium gluconate 1 g IV over 3 minutes. Repeat doses may be necessary. Calcium chloride can also be used in lieu of calcium gluconate. The suggested dose for calcium chloride for magnesium toxicity is 500 mg of 10% calcium chloride IV given over 5-10 minutes.
5. In the setting of severe preeclampsia, magnesium sulfate should be administered upon diagnosis and continued until 24 hours after birth.
 - a. However, if the patient shows no improvement in her symptoms of preeclampsia, clinical judgment is advised and magnesium sulfate administration may need to be provided for an extended period of time.
 - b. If the patient is stabilized and remote from term and being expectantly managed, we advise continuing magnesium sulfate during the course of administering corticosteroids for fetal lung maturity. Upon completion of administration of the corticosteroids, magnesium sulfate can be stopped. Magnesium sulfate should be resumed if there are escalating signs of worsening preeclampsia, eclampsia, or plans are made to proceed with delivery.
6. In the setting of mild preeclampsia the use of magnesium sulfate for seizure prophylaxis can be considered.
7. If an eclamptic seizure occurs postpartum and the patient is off of magnesium sulfate, it should be re-administered for at least 24 hours after the last seizure.
8. If a patient has recurrent seizures despite already being on magnesium sulfate the first therapy we recommend is an additional loading dose of magnesium sulfate 2g IV over 5 minutes. If the patient continues to have seizures despite a repeat loading dose of magnesium sulfate, alternative anti-convulsants should be considered:
 - a. Lorazepam (Ativan) 4mg IV over 2-5 minutes (can repeat in 5-15 minutes) to maximum of 8 mg in 12 hours
 - b. Diazepam (Valium) 5-10mg IV slowly (can repeat every 15 minutes up to 30mg)
 - c. Midazolam (Versed) 1-2mg IV (can repeat in 5 -10 minutes)
 - d. Phenytoin (Dilantin) 1000 mg IV over 20 minutes

- e. MLD--Use of other agents such as Propofol will be institution specific.
9. Consideration of severe CNS events must be entertained for recurrent or persistent seizures.
 10. Renal insufficiency: magnesium sulfate should be used with caution in women with renal insufficiency/failure (i.e. serum creatinine greater than 1.2 mg/dL). In these patients, if they are naive to magnesium therapy, a loading dose can be administered. A lower maintenance dose can be considered with serial serum magnesium levels to guide therapy (e.g. 1 g per hour).
 11. Myasthenia Gravis: magnesium sulfate is contraindicated in patients with myasthenia gravis.

EVIDENCE GRADING


Level of Evidence: I-A, I-B, II-B, III, IV

REFERENCES

1. Magnesium Sulfate: Drug Information Online, updated 4/2013. 2012; <http://www.drugs.com/pro/magnesium-sulfate.html>. Accessed 6/14, 2013.
2. Altman D, Carroli G, Duley L, et al. Do women with pre-eclampsia, and their babies, benefit from magnesium sulphate? The Magpie Trial: a randomised placebo-controlled trial. *Lancet*. 2002;1(359(9321)):1877-1890.
3. Coetzee E, Dommissie J, Anthony J. A randomised controlled trial of intravenous magnesium sulphate versus placebo in the management of women with severe pre-eclampsia. *Br J Obstet Gynaecol*. 1998;105(3):300-303.
4. Duley L, Gulmezoglu A, Chou D. Magnesium sulphate versus lytic cocktail for eclampsia. In: 2010;8(9):CD002960 CDSR, ed2010.
5. Duley L, Henderson-Smart D. Magnesium sulphate versus phenytoin for eclampsia. In: 2003;4(4):CD000128 CDSR, ed2003.
6. Duley L, Matar H, Almerie MQ, Hall D. Alternative magnesium sulphate regimens for women with preeclampsia and eclampsia. In: 2010;4(8):CD007388 CDSR, ed2010.
7. Duley L, Matar HE, Almerie MQ, Hall DR. Alternative magnesium sulphate regimens for women with pre-eclampsia and eclampsia. *Cochrane Database of Systematic Reviews*. 2010(8):CD007388.
8. Ehrenberg H, Mercer B. Abbreviated postpartum magnesium sulfate therapy for women with mild preeclampsia: a randomized controlled trial. *Obstet Gynecol*. 2006;108(4):833-838.

9. Livingston J, Livingston L, Ramsey R, Mabie B, Sibai B. Magnesium sulfate in women with mild preeclampsia: a randomized controlled trial. *Obstet Gynecol.* 2003;101(2):217-220.
10. Alexander J, McIntire D, Leveno K, Cunningham F. Selective magnesium sulfate prophylaxis for the prevention of eclampsia in women with gestational hypertension. *Obstet Gynecol.* 2006;108(4):826-832.
11. ACOG. Diagnosis and Management of Preeclampsia and Eclampsia #33. *American College of Obstetricians and Gynecologists Practice Bulletin Number 33.* 2002 (Reaffirmed 2008).
12. World Health Organization. WHO Recommendations for Prevention and Treatment of Preeclampsia and Eclampsia: Evidence Base. 2011.
13. Magee LA, Abalos E, von Dadelszen P, Sibai B, Easterling T, Walkinshaw S. How to manage hypertension in pregnancy effectively. *Br J Clin Pharmacol.* Sep 2011;72(3):394-401.
14. Visintin C, Mugglestone M, Almerie MQ, Nherera L, James DC, Walkinshaw S. Management of hypertensive disorders during pregnancy: summary of NICE guidance. *BMJ.* 2010;25(341(25)):c2207.
15. Beyea S, Kobokovich L, Becker S, Hicks R. Medication Errors in the LDRP: Identifying Common Errors Through MEDMARX Reporting. *AWHONN: Lifelines.* 2004;8(2):130-140.
16. Leveno K, Alexander J, McIntire D, Lucas M. Does magnesium sulfate given for prevention of eclampsia affect the outcome of labor? *Am J Obstet Gynecol.* 1998;178(4):707-712.
17. Atkinson M, Guinn D, Owen J, Hauth J. Does magnesium sulfate affect the length of labor induction in women with pregnancy-associated hypertension? *Am J Obstet Gynecol.* 1995;173(4):1219-1222.

ECLAMPSIA ALGORITHM



SAMPLE PREECLAMPSIA/ECLAMPSIA MEDICATION BOX

(Each institution should prepare its own medication box)

Contents of Labor & Delivery Severe Pre-Eclampsia & Eclampsia Box	
Drug contents	
Medications	Quantity
Magnesium sulfate 20g/500ml bag	1
Calcium gluconate 1000mg/10ml vial	1
Labetalol 100mg/20ml vial	1
Hydralazine 20mg/ml vial	1
Esmolol 100mg/10ml vial	1
Sodium Chloride 0.9% (NS) 10 ml vial	1
Propofol 10mg/ml, 20ml vial	
Supply contents	
Supplies	Quantity
20 ml syringe	1
10 ml syringe	2
3 ml syringe	2
18G needle (pink)	5
Alaris Pump Tubing Set	1

Kindly used with permission of Stanford University Medical Center and Gillian Hilton, MD, 2013.

TEAMWORK AND COMMUNICATION

Audrey Lyndon, PhD, RNC, FAAN

Linda Walsh, CNM, MPH, PhD, FACNM

Racine Edwards-Silva, MD, FACOG Olive View UCLA Medical Center

Effective communication and teamwork are essential components of obstetric safety and quality.¹ Communication breakdowns and failures of organizational culture and teamwork have consistently ranked among the top three leading contributors to reported maternal and newborn sentinel events.^{2,3} While it is difficult to link specific, discreet communication strategies to changes in patient outcomes, there is evidence that sustained attention to communication, teamwork, and safety can indeed improve perinatal outcomes.⁴⁻⁶ Furthermore, empowerment of staff to speak up when they see problems or lack of protocol compliance has been a central component of initiatives to reduce or eliminate complications such as central line infections that were previously deemed unpreventable, and teamwork training substantially reduced surgical mortality.⁷⁻¹⁰ However, multiple studies suggest that clinicians frequently may remain silent about their clinical concerns even when they know or believe that harm could potentially result from continuing with planned care.¹¹⁻¹⁵ Patients can also identify clinical deterioration and lack of protocol compliance, but often feel ignored if and when they raise concerns.¹⁶⁻¹⁸ The seven U.S. professional organizations for clinicians who care for childbearing women assert that shared decision making, effective communication, and effective teamwork are fundamental tenets of quality patient care.¹ These principles are especially important in the setting of complications such as preeclampsia/eclampsia, where the potential for catastrophic problems is elevated and early identification and communication of disease progression is essential for effective management.¹⁷

Preeclampsia/eclampsia was the second leading cause of maternal deaths in the 2002-2003 California Pregnancy-Related Mortality Review.¹⁹ For women whose deaths were reviewed, clinician factors were identified as contributory to the death in 78% of cases, and facility or system level factors were identified as contributory to the deaths in 57% of cases. Overall, 48% of preeclampsia/eclampsia deaths were determined to have a strong or good chance to alter outcomes. Delays or failures in treatment, misdiagnosis, and denial of the severity of women's illness were key factors contributing to fatal outcomes.¹⁹ Failures in communication, including silence in the face of clinical concern and lack of listening skills or responsiveness to concern, are likely to have contributed to delays, misdiagnosis, and treatment failures. Therefore, creating an environment where all staff—regardless of formal or informal status within the medical and organizational hierarchy—can and will speak up about their clinical concerns is critical to implementing clinical practice changes to improve preeclampsia outcomes and clinical safety.

Effective, highly reliable teams are preoccupied with the potential for failure and therefore collectively monitor and crosscheck each other and clinical processes to proactively

identify potential problems. It is essential that perinatal units create an environment where all staff are empowered to “stop the line,” or formally interrupt planned care and procedures to check safety when they observe there is potential for harm.^{20,21} Key skills include the ability of all staff, patients, and family members to speak up about concerns with persistence until a mutually agreeable resolution is established, and the ability to listen to each other and respond in a supportive manner regardless of whether or not they agree with their peers.^{1,22} Established strategies for improving communication and teamwork are well delineated nationally and internationally, and are outlined in Table 1.^{17,23}

Perspectives on the best course of action for a woman with preeclampsia in specific clinical situations may vary between physicians, midwives, nurses, the woman, and the woman’s family members (see Table 2). Managing conflict traditionally has been difficult for clinicians. Key skills for handling conflict effectively include: a) addressing it rather than letting concerns fester; b) taking the time to listen carefully to the concerns of others; c) setting aside assumptions, especially regarding what motivates others’ behavior; d) being willing to own part of the problem. When clinical disagreements are approached with a spirit of inquiry, good will, active listening, and dedication to shared decision-making, they can often be resolved quickly and in a manner that builds continued trust between team members.²² In the event that concerns cannot be resolved using these or other communication strategies, all clinicians, including registered nurses, have an affirmative duty to pursue their concerns through the institutional chain of authority (Figure 1).²⁴

Culture shifts including communication and behavior change require commitment from all staff across all disciplines. Numerous training models are available to assist in assessment, development, implementation and evaluation of communication and collaboration in complex settings. Findings of interdisciplinary team training have suggested that focused training contributes to optimizing human performance and reducing human error.¹⁷

RECOMMENDATIONS FOR QUALITY IMPROVEMENT

Recommendation Strength B-C

- 1) All units should adopt the principles outlined in “Quality Patient Care in Labor and Delivery: A Call to Action”¹
- 2) Adoption of standardized protocols supports effective teamwork. Units should standardize protocols for risk assessment, medication selection and administration, and parameters for patient monitoring and primary provider notification.
- 3) Policy, procedure, and unit culture should outline clear lines of communication and clear avenues for escalation when appropriate.
- 4) Administrators must support clinicians and patient who raise concerns.

5) Conduct routine briefings and debriefings for patients with preeclampsia.

Table 1. Communication Strategies to Foster Mutual Respect and Shared Decision-making (30)

Briefings	<ul style="list-style-type: none"> • Set tone for team interaction • Can be a routine part of board rounds, huddles, handouts and bedside rounds
Debriefings	<ul style="list-style-type: none"> • Used to identify what happened, what was learned, and what can be done better next time • Can be team-building in real patient situations as well as simulation learning
Assertive Language	<ul style="list-style-type: none"> • Effective assertion is persistent, polite, timely, clear and solution focused • Using “CUS” as a guideline: “I’m Concerned,” “I’m Uncomfortable”, “this is a Safety Issue”
Critical Language	<ul style="list-style-type: none"> • Ensures that specific, relevant, critical information is communicated; example: SBAR (Situation, Background, Assessment, Recommendation)
Closed Communication Loop	<ul style="list-style-type: none"> • Receiver of information restates what was said to the sender to ensure correct understanding. • Reinforces the importance of effective listening
Call Outs	<ul style="list-style-type: none"> • Used to confirm the phase of a process

Adapted from: Teamwork and Communication Working Group. Improving patient safety with effective teamwork and communication: Literature review needs assessment, evaluation of training tools and expert consultations, 2011. Edmonton (AB): Canadian Patient Safety Institute; TeamSTEPPS®: Strategies and Tools to Enhance Performance and Patient Safety, Agency for Healthcare Research and Quality, and Quality Patient Care in Labor and Delivery: A Call to Action. *J Midwifery Women’s Health*. 2012; 57(2):112-113 and *J Obstet, Gynecol Neonatal Nurs*. 2012;41(1):151-3.

Example dialogue—Closed Communication Loop

Nurse: Hi CNM Jones, this is Nurse Smith, calling about Ms. Green in room 27 at XYZ birth center. She is 39-3/7 weeks G1P0 admitted for nausea and vomiting this morning. Her blood pressure is 150/92, no proteinuria on dip UA, but she has a sudden severe headache. I am concerned and would like you to come over now to evaluate her.

CNM: Thanks Nurse Smith, I’m going to be over later to rupture her membranes and get this labor going. She must be miserable from all that vomiting and probably has the flu.

Nurse: Hmm. I understand the vomiting could be a GI bug. But I’m concerned that the signs and symptoms Ms. Green is demonstrating could also be atypical preeclampsia, and if so, the headache would make it severe preeclampsia. I really think she needs a workup now and you should come over to evaluate her. When can I expect to see you?

CNM: Oh, I see. Please draw xyz labs right away. I'll be right over, and I'm calling the OB back-up now. Thanks for clarifying your concerns.

Nurse: Ok great. I'll draw x, y, and z labs right away. I'll let Ms. Green know you'll be in to see her in about 15 minutes.

CNM: Agreed, thank you.

Draft

Table 2. Approaches for improving communication and resolving clinical disagreements

Sources of Potential Conflict	Approach – May Need to have:
Differing expectations for information needs, communication content and style	<ul style="list-style-type: none"> • Team Training • Structured communication tools (e.g., SBAR-R-R structured handoffs)^a; • Board rounds • Huddles • Attentive listening
Failure to communicate rationale Inattention to concern Concerns remain unresolved	<ul style="list-style-type: none"> • Routinely ask for plan and reasoning • Persistently restate concerns until resolved • Develop clear lines for problem resolution that can be activated quickly with high risk patients: e.g. laborist in house; MFM consultation available 24 hours a day; back-up list for who to call including anesthesiologists, MFM, intensivists, and administrators • Ratify plan before concluding conversation
Differing “world views,” e.g., use of magnesium in women with mild preeclampsia; meaning of signs and symptoms such as nausea, lethargy, or headache; interpretation, and management of complex tracings	<ul style="list-style-type: none"> • Standardize protocol for magnesium sulfate, including criteria for administration • Standardize ongoing clinical assessments and notification parameters for signs of potential disease progression or magnesium toxicity • Standardize fetal monitoring language and application • Provide regular interprofessional case reviews to discuss management; role model expression of concern and positive resolution of differences • Standardize expectations for notification of complications • Articulate and plan for potential problems early in care • Individuals take responsibility for collaboratively discussing differing views • Avoid professional stereotyping as an explanation for behavior
Disruptive behavior	<ul style="list-style-type: none"> • “Good Citizen” policy consistently enforced • Individuals and peers stand up to unprofessional behaviors • Administrative commitment to addressing any chronic issues • Availability of anonymous incident reporting system

Adapted from Lyndon, Zlatnik & Wachter Effective physician-nurse communication: a patient safety essential for labor and delivery. *Am J Obstet Gynecol.* 2011;Aug;205(2):91-6.

^aSBAR-R-R = Situation, Background, Assessment, Recommendation, Reasoning, Ratification

SBAR-R-R Communication Technique

A specific strategy for structured communication that many health care providers are familiar with is “SBAR.” This format, developed by Kaiser Permanente, was adapted from military and aviation crew resource management practices. It is recommended and taught in most healthcare teamwork improvement programs. The SBAR format, which stands for Situation-Background-Assessment-Recommendation, provides a brief, organized, predictable flow of information that facilitates critical thinking and communication skills between healthcare providers, and may be especially helpful in leveling communication styles between disciplines. However SBAR alone does not explicitly incorporate essential teamwork principles of assertive communication of concern and closed loop communication. These two principles can be built into SBAR with a simple expansion to SBAR-R-R (Table 3), which includes the steps “Reasoning,” to ensure team members understand each other’s interpretation of the present situation if immediate agreement is not reached, and “Ratification,” to ensure the team members have an agreed upon plan for moving forward.

Table 3. SBAR-R-R Communication Technique Applied to Preeclampsia

Prepare for an SBAR-R-R by:

1. Assessing the patient
2. Reviewing recent notes and laboratory results
3. Having the medical record available during the conversation

Situation: Always identify yourself, where you are calling from, the name of the woman you are calling about, quickly state the main reason and the level of urgency for the call.

Background: Give brief pertinent background information – medical history, complaints, vital signs, and interventions that have already occurred

Assessment: Say what you think is going on

Recommendation: Say what you think should happen or ask for specific orders

Reasoning: If the response is not what you expect and requested, state why what you think should happen is important. What could happen if we don’t do this?

Ratification: Close the loop by confirming actions to be taken. Assure mutual agreement on the plan.

	Ambulatory Care or Emergency Department	Inpatient Antepartum or Intrapartum	Postpartum
Situation	<p>I am calling about Ms. ____, who</p> <ul style="list-style-type: none"> <input type="checkbox"/> is pregnant <input type="checkbox"/> recently had a baby <p>and is here in the ED with stomach pain. I am concerned about</p> <ul style="list-style-type: none"> • High blood pressure • Headache • Visual disturbances • Decreased fetal movement • Nausea and vomiting 	<p>I am calling about Ms. ____, who is an antepartum patient being monitored for preeclampsia. I am concerned about:</p> <ul style="list-style-type: none"> • New onset headache • Increasing blood pressures • Headache that has not resolved • Visual disturbances • Stomach pain • Abnormal or indeterminate fetal status • Altered/worsening lab values 	<p>I'm calling about Ms. ____ who had her second baby yesterday at 3 pm. I am concerned about:</p> <ul style="list-style-type: none"> • New onset headache • Increasing blood pressures • Headache that has not resolved • Visual disturbances • Stomach pain • Altered/worsening lab values
Background	<ul style="list-style-type: none"> • GPTAL @__weeks or G_P_ #days post birth • Significant OB and medical history • Current problems • Patient complaints • Vital Signs • Interventions and response 	<ul style="list-style-type: none"> • GPTAL @__weeks • Significant OB and medical history • Current problems • Patient complaints • Vital Signs • FHR tracing baseline, variability, accelerations, decelerations • Uterine activity • Interventions already completed 	<ul style="list-style-type: none"> • G_P__ • Mode of birth (vaginal/cesarean) • Significant OB and medical history • Current problems • Patient complaints • Vital Signs • Interventions already completed
Assessment	<ul style="list-style-type: none"> • I'm thinking she may have preeclampsia and need an OB evaluation before we can clear her. • I'm concerned she may have severe preeclampsia and needs medication to 	<ul style="list-style-type: none"> • Her preeclampsia seems to be progressing and her blood pressures indicate severe hypertension and severe preeclampsia. • The FHR tracing is 	<ul style="list-style-type: none"> • I'm thinking that her increasing BPs and new onset headache may represent preeclampsia and that she would benefit from an initial preeclampsia workup.

	control her blood pressure now.	indeterminate and the decelerations do not resolve with position change.	
Recommendation	<ul style="list-style-type: none"> • Could you please come and evaluate her within _____? <ul style="list-style-type: none"> ○ Now ○ Within 30 min ○ Before____, etc. • Could I have orders for:_____ <ul style="list-style-type: none"> ○ CBC, liver function, kidney function ○ Antihypertensive • Magnesium sulfate 	<ul style="list-style-type: none"> • I need you to come and evaluate her now. • May I please have an order for antihypertensive medication? • Are there any labs we need to repeat? • When can I expect you? 	<ul style="list-style-type: none"> • May I have an order for a preeclampsia lab panel? • When can I expect you in to evaluate Ms. ____?
Reasoning	<ul style="list-style-type: none"> • I don't think it is safe to send her home without evaluating the possibility of preeclampsia • If we don't lower her blood pressure to a safer range she could have a stroke 	<ul style="list-style-type: none"> • It is really important to control her blood pressure while we make preparations to proceed to birth. • If we don't lower her blood pressure to a safer range she could have a stroke. 	<ul style="list-style-type: none"> • It's important for us to get baseline data before considering discharge in the morning.
Ratification	<ul style="list-style-type: none"> • Ok, I'll do _____, and You'll evaluate her in _____ or call _____ for _____. 	<ul style="list-style-type: none"> • Ok, I'll do _____, and you'll be here to evaluate her in _____. 	<ul style="list-style-type: none"> • OK, I'll do ____ and you'll be in to evaluate her in _____.

Adapted from Kaiser Permanente SBAR Guidelines and SBAR Report to Physician about a Critical Situation, and Ascension Health Perinatal SBAR Report Template.

Example Strategy for Building Collaborative Culture and Problem Solving Skills
MOPS (Nurse-Led Multidisciplinary Obstetric Patient Summaries)

Every patient is discussed by the multidisciplinary team each shift. This might occur at board rounds with the entire L&D team, or might be a two-person process involving the attending physician or midwife and bedside RN, with additional consultation from anesthesia, MFM, charge nurse, or others as needed for patient complexity. The exact make-up and logistics for each team are dependent on local conditions.

All care providers are encouraged to consider elements of concern or potential risks by pondering questions such as:

- What potential risks exist for this patient? (Is there risk of stroke, eclampsia, hemorrhage, fetal injury?)
- Are there trends that indicate concern? (e.g., vital signs, fetal trends, lab trends, headache, malaise, nausea, abdominal pain, scotomata)
- Is there any information or task that I don't understand or know how to perform?
- What is the plan of care based on the given information?
- Do I feel uncomfortable or I am concerned about the plan of care?
- Do I feel qualified or do I feel inexperienced in caring for a patient like this?
- Are there concerns I would like to have addressed?

EVIDENCE GRADING:


Level of Evidence: II-3, III

REFERENCES

1. Quality Patient Care in Labor and Delivery: A Call to Action. *J Midwifery Women's Health*. 2012;57(2):112-113.
2. The Joint Commission. Preventing infant death and injury during delivery. Sentinel Event Alert. 2004;Jul 21(30):1-3.
3. The Joint Commission. Sentinel Event Data Root Causes by Event Type 2004-2012. 2012;
http://www.jointcommission.org/assets/1/18/Root_Causes_Event_Type_2004_2Q2012.pdf.
4. Pettker CM, Thung SF, Norwitz ER, et al. Impact of a comprehensive patient safety strategy on obstetric adverse events. *Am J Obstet Gynecol*. Feb 26 2009.
5. Pettker C, Thung S, Raab C, et al. A comprehensive obstetrics patient safety program improves safety climate and culture. *Am J Obstet Gynecol*. 2011;204(3216.e1-e6).
6. Simpson K, Knox G, Martin M, et al. Michigan Health & Hospital Association Keystone Obstetrics: a statewide collaborative for perinatal patient safety in Michigan. *Joint Commission Resources*. 2011;Dec;37(12):544-552.
7. Miller M, Niedner M, Huskins W, et al. Reducing PICU central line-associated bloodstream infections: 3-year results. *Pediatrics*. 2011;Nov(128(5)):e1077-1083.
8. Lin D, Weeks K, Bauer L, et al. Eradicating central line-associated bloodstream infections statewide: the Hawaii experience. *Am J Med Qual*. 2012;27(2):124-129.
9. Dreifus C. A Conversation with Dr. Peter J. Pronovost: Doctor Leads Quest for Safer Ways to Care for Patients. *New York Times* 2010: Sec D2.
10. Neily J, Mills P, Young-Xu Y, et al. Association between implementation of a medical team training program and surgical mortality. *JAMA*. 2010;Oct 20(304(15)):1693-1700.
11. Blatt R, Christianson M, Sutcliff K, et al. A sensemaking lens on reliability. *Journal of Organizational Behavior*. 2006;27:897-917.

12. Lyndon A. Social and environmental conditions creating fluctuating agency for safety in two urban academic birth centers. *J Obstet Gynecol Neonatal Nurs*. 2008;Jan-Feb(37(1)):13-23.
13. Smetzer J, Cohen M. Intimidation: practitioners speak up about this unresolved problem. Joint Commission Journal on Quality and Patient Safety. *Journal on Quality and Patient Safety*. 2005;31:594-599.
14. Lyndon A, Sexton JB, Simpson KR, Rosenstein A, Lee KA, Wachter RM. Predictors of likelihood of speaking up about safety concerns in labour and delivery. *BMJ Qual Saf*. Jul 1 2011.
15. Gardezi F, Lorelei L, Sherry E, et al. Silence, power and communication in the operating room. *J Adv Nurs*. 2009;65(7):1390-1399.
16. Entwistle V, Mello M, Brennan T. Advising patients about patient safety: current initiatives risk shifting responsibility. *Jt Comm J Qual Patient Saf*. 2005;Sep(31(9)):483-494.
17. Teamwork and Communication Working Group. Improving patient safety with effective teamwork and communication: Literature review needs assessment, evaluation of training tools and expert consultations. 2011; <http://www.patientsafetyinstitute.ca/English/toolsResources/teamworkCommunication/Documents/Canadian Framework for Teamwork and Communications.pdf>.
18. Schwappach D. Review: Engaging Patients as Vigilant Partners in Safety. *Med Care Res Rev*. 2010;67(2)(April 1):119-148.
19. *The California Pregnancy-Associated Mortality Review. Report from 2002 and 2003 Maternal Death Reviews: California Department of Public Health, Maternal Child and Adolescent Health Division*. Sacramento 2011.
20. Leonard M, Graham S, Bonacum D. The human factor: the critical importance of effective teamwork and communication in providing safe care. *Qual Saf Health Care*. 2004;Nov(13):Sup 1:885-890.
21. Knox G, Simpson K. Perinatal high reliability. *Am J Obstet Gynecol*. 2011;May(204(5)):373-377L.
22. Lyndon A, Zlatnik MG, Wachter RM. Effective physician-nurse communication: a patient safety essential for labor and delivery. *American Journal of Obstetrics and Gynecology*. Aug 2011;205(2):91-96.
23. AHRQ. TeamSTEPPS®: Strategies and Tools to Enhance Performance and Patient Safety. 2012; <http://teamstepps.ahrq.gov/abouttoolsmaterials.htm>.
24. American Nurses Association. Code of Ethics for Nurses with Interpretive Statements. *American Nurses Association 2001*.

Example of resources for collaborative management of clinical disagreement and escalation cycle for conflict resolution



THE ROLE OF MEDICAL SIMULATION

Mark Meyer, MD, Kaiser Permanente San Diego

BACKGROUND

The use of medical simulation and the recognition of its value continues to grow. There is a growing body of literature that demonstrates the efficacy of medical simulation, but a review of this data is beyond the scope of this discussion.

It is important to recognize that medical simulation represents a spectrum of tools that spans from low fidelity drills to high fidelity, inter-professional, interdisciplinary team simulations. Effective simulation programs must be designed with clear learning objectives and tailored to available resources and instructor expertise. They do not require extensive resources to be effective.

While simulation can be used to address knowledge gaps in the identification and treatment of preeclampsia, its greatest value lies in its potential to help teams “put together.” Whether conducted in a dedicated simulation lab or in real patient care (in-situ simulation), inter-professional team training allows for:

- Testing of new policies and procedures
- Demonstration of skills in a more realistic environment
- Identification of systems issues and the ability to test new systems
- Instruction in techniques to improve communication and coordination of treatment teams e.g., human factors, etc.

EDUCATIONAL MATERIALS

When constructing simulations for preeclampsia and eclampsia there are several variables to consider:

1. Different locations:
 - Location of the patient –LDR (labor and delivery room), Postpartum, L&D triage, Emergency Department
2. Different diagnoses:
 - Diagnosis: mild vs. severe preeclampsia, eclampsia,
 - Requirements for successful treatment – e.g. will one (1) dose of anti-hypertensive medication be sufficient, or will the team be required to proceed with multiple doses/drugs to be successful?
 - Simulation expertise and resources of the participating medical center.

It is not possible to create a scenario that addresses all these variables. For the purposes of this program, two sample scenarios are provided that require the management of severe preeclampsia and eclamptic seizures that are refractory to magnesium. Both scenarios provide detailed information essential for an effective simulation experience, including debriefing guides specific to the scenario and a generic debriefing guide for teamwork/communication skills.

Scenarios provided include:

1. A version that covers both low fidelity and high fidelity simulations.
2. A detailed version created for teams that have high fidelity capabilities. This particular version includes programming and details for teams using SimMan3G.

There is no expectation that these scenarios alone will be adequate. Rather, these are simply examples that can be modified to include different clinical environments, differing levels of clinical complexity, and differing levels of simulation expertise/resources.

The scenarios are listed below and can be found in the appendices G through N.

- Severe Preeclampsia and Eclampsia in LDR v2.0 SimMan3G
 - Scenario Part 1: General Information
 - Scenario Part 2: Learning Objectives
 - Scenario Part 3: Patient Background Information
 - Scenario Part 4: Equipment/Materials List
 - Scenario Part 5: Program Algorithm & GUI (Graphic User Interface) Notes
 - Scenario Part 6: Debriefing Objectives
 - Scenario Part 7: Debriefing Guide/Evaluation

Simulations can be done with or without SimMan3 equipment.

FLUID MANAGEMENT IN PREECLAMPSIA

Tom Archer, MD, MBA, University of California San Diego
Holly Champagne, MSN, RNC, CNS, Kaiser Permanente Northern California

BACKGROUND

Fluid management in preeclampsia is often difficult because of a leakage of water, electrolytes, and plasma from the intravascular space, due to underlying endothelial damage. This leakage can produce significant fluid shifts into the interstitial space resulting in peripheral and/or central (pulmonary and central nervous system, CNS) edema. As fluid shifts out of the intravascular space, there is also the potential for hypovolemia. Therefore, fluid administration must be assessed in the context of preserving organ perfusion, while limiting or preventing pulmonary edema. Renal endothelial damage appears to be particularly sensitive to these fluid changes resulting in proteinuria and oliguria. Assessment of renal function (serum creatinine) should be assessed to determine the degree of renal dysfunction. One hallmark of intravascular depletion is hemoconcentration.¹⁻⁴ Since pulmonary edema is more common and permanent renal damage due to preeclampsia is rare, fluids are normally restricted (see section related Magnesium administration).⁵

If oliguria occurs, a trial of intravenous (IV) fluid bolus of 250-500 ml isotonic fluid (normal saline or Lactated Ringer's) can be given. If after a total infusion of 1000 mL of crystalloid IV fluids oliguria is not resolved, consideration is usually given to other modalities for enhancing urine output. If hypovolemia is still suspected then a trial of colloid administration or blood may be used. Oxygen saturation monitoring is essential in these clinical settings due to the risk of pulmonary edema with excessive fluid administration in preeclamptic patients. If the patient is felt to be adequately hydrated, a trial pharmacological diuresis (furosemide, 5-10 mg IV) may be attempted. Oliguria, particularly when combined with excessive IV fluid administration, significantly increases the risk for pulmonary edema. Oliguria will also reduce the renal clearance of magnesium and the maintenance dose of magnesium will need to be adjusted to reduce the risk of magnesium toxicity. In the setting of oliguria and reduced O₂ saturation (i.e. below 95%) diuresis is indicated. In rare instances, placement of a pulmonary artery catheter and measurement of pulmonary capillary wedge pressure may be needed. If these steps become necessary, the patient should be transferred to the intensive care unit for hemodynamic monitoring

KEY LEARNING POINTS

1. Patients with severe preeclampsia should have strict fluid intake and output monitoring assessments.

2. Total fluid intake (oral and intravenous) should be limited in both mild and severe preeclampsia. Many recommend that the sum of oral and all IV fluid should be ≤ 125 mL / hour (range 60 to ≤ 125 mL/hr) unless there are other clinical circumstances that dictate a different management plan.
3. Serum creatinine should be assessed in all patients with gestational hypertension, preeclampsia, or chronic hypertension with superimposed preeclampsia.
4. A Foley catheter with urometer is useful for monitoring urine output and is essential in the setting of oliguria or pulmonary edema.
5. An oliguric patient (less than 30 mL per hour for two hours, or less than 500 mL in 24 hours), should be given a limited trial of IV fluid boluses (normal saline or Lactated Ringer's), usually starting with 250-500 mL.
6. After a total of 1000 mL of crystalloid IV fluid is administered without resolution of the oliguria, and hypovolemia is still suspected, consideration may be given to the administration of colloid (e.g., albumin) or blood to enhance renal perfusion and urine output.
7. After adequate hydration, consideration should be given to the use of pharmacological diuresis (furosemide).
8. In the setting of oliguria and reduced O₂ saturation (i.e., below 95%), pulmonary edema should be strongly suspected and diuresis is indicated.
9. Anesthesiologists should consider omitting or reducing the fluid bolus prior to epidural analgesia and ensure proper lateral positioning to avoid hypotension. Early treatment of hypertension has consistently been found to reduce the incidence of hypertensive crisis and will decrease the risk or prevent intracranial hemorrhage.^{2-4,6}
10. Cardiac dysfunction should be strongly considered in the presence of persistently low O₂ saturations, the development of respiratory distress, persistent pulmonary edema, or unanticipated low BP's. In this setting, measurement of maternal BNP (Brain Natriuretic Peptide) and/or maternal echocardiography is strongly recommended to evaluate cardiac function and detect dilated cardiomyopathy.

EVIDENCE GRADING

Level of Evidence: IIIC

REFERENCES

1. Aukes A, Wessel I, Dubois A, Aarnoudse J, Zeeman G. Self-reported cognitive functioning in formerly eclamptic women. *Am J Obstet Gynecol*. 2007;197:365.e361--365.e366.
2. Postma I, Wessel I, Aarnoudse JG, Zeeman G. Neurocognitive functioning in women with a history of eclampsia: executive functioning and sustained attention. *American Journal of Perinatology*. 2010;27(9):685-690.
3. Wiegman M, de Groot J, Jansonius N, et al. Long-Term Visual Functioning After Eclampsia. *Obstet & Gynecology*. 2012;119(5):959-966.
4. Zeeman G, McIntire D, Twickler D. Maternal and fetal artery Doppler findings in women with chronic hypertension who subsequently develop superimposed pre-eclampsia. *J Matern Fetal Neonatal Med*. 2003;Nov(14(5)):318-323.
5. Creasy R, Resnik R, Iams J, Lockwood C, Moore T. *Creasy & Resnik's Maternal-Fetal Medicine: Principles and Practice*. Vol Sixth Edition, Chapter 35, pg. 673. 2008.
6. Creasy R, Resnik R, Iams J, Lockwood C, T M. *Creasy & Resnick's Maternal Fetal Medicine: Principles and Practice*. Vol Sixth Edition, Chapter 35, pg 673-674. 2008.

AIRWAY MANAGEMENT IN PREGNANT OR POSTPARTUM WOMEN HAVING SEIZURES

Tom Archer, MD, MBA, University of California San Diego

BACKGROUND

A seizure is a frightening and uncommon occurrence in Labor and Delivery and the visceral response of many providers is to immediately administer magnesium sulfate to stop the abnormal movement associated with the seizure. However, even more important than stopping the seizure (which usually stops on its own after 1-2 minutes) is maintaining and protecting the airway. Seizures do not directly cause death, but intracranial hemorrhage and hypertensive encephalopathy do.^{1 2} The airway is the first priority in seizure management, even before administration of magnesium sulfate.

Basic airway maintenance skills need to be re-taught and actively maintained by nurses and physicians who work in Labor and Delivery, since this “skill set” is used very infrequently in that environment. Anesthesiologists are accustomed to caring for unconscious patients who may not be breathing adequately. For this reason, the anesthesiologist should be called immediately when a patient suffers a seizure and an anesthesiologist should be involved in teaching the Labor and Delivery staff about airway management.^{1,2}

KEY LEARNING POINTS

1. Seizures involve loss of consciousness and violent movements, with the potential for blocked ventilation (“airway obstruction”), regurgitation and aspiration of gastric contents, as well as falls, head trauma and tongue biting.
2. Many of the sequelae of seizures, such as hypertensive brain hemorrhage and fetal and maternal hypoxic brain damage, are due to the cessation of respiration and the failure to deliver oxygen to the maternal brain and the placenta.
3. Maternal hypoxia is the most common cause of “fetal distress” following an eclamptic seizure.
4. Anesthetizing the preeclamptic mother may present special challenges related to lack of maternal cooperation for neuraxial anesthesia and for obese patients, swelling of the airway tissues that makes intubation more difficult. Therefore, emergency induction of anesthesia may lead to both maternal and fetal compromise.

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. Call for help. Notify the anesthesiologist immediately.
2. Turn patient into a lateral 'recumbent' position. The side-lying position prevents aortocaval compression, helps the tongue fall to the side of the mouth and lessens the risk of aspiration. If possible cushion the head from injury by placing a soft object under the head.
3. Open airway with a jaw thrust and/or oral airway, if needed. Do not insert any object other than the oral airway, if needed, into the person's mouth. Be aware that nasal airways will often cause nosebleed. Check for air movement and reposition if there is no air movement. Be aware that an oral airway can make the patient vomit and may not be necessary.
4. Apply oxygen, obtain suction, and pulse oximeter to check SPO₂.
5. Obtain IV access.
6. Administer magnesium sulfate. (Refer to Magnesium Sulfate chapter, pg. 48)
7. Control blood pressure if necessary with IV meds, but be aware that hypoxia and hypercarbia will elevate blood pressure.
8. The initial focus should be on opening and protecting the airway and supplying the patient with oxygen. Laryngoscopy will result in an acute hypertensive episode, so pre-intubation medication such as IV Esmolol, IV Lidocaine, or IV Remifentanyl should be administered.
9. Resuscitation of the mother is the key to protecting the fetus. This point is counterintuitive for many Labor and Delivery personnel, who may incorrectly focus on the baby, when it is the mother who requires top priority during and after a seizure.
10. Following a maternal seizure, fetal bradycardia is commonly seen due to maternal hypoxia. Stabilization of the mother is the first priority followed by resuscitation of the fetus. Cesarean section should be reserved for the situation in which maternal and fetal resuscitation are unsuccessful in stabilizing the mother or resolving the non-reassuring fetal heart rate tracing.
11. Regular drills should be conducted in Labor and Delivery for management of the airway during an eclamptic seizure.

EVIDENCE GRADING
Level of Evidence: III-C

REFERENCES

1. Creasy R, Resnik R, Iams J, Lockwood C, Moore T. *Creasy & Resnik's Maternal-Fetal Medicine: Principles and Practice*. Vol Sixth Edition, Chapter 35, pg. 673. 2008.
2. Chestnut D, Polley L, Tsen L, Wong C. *Chestnut's Obstetric Anesthesia: Principles and Practice, 4th Edition, Chapter 45, pg 984-986*. 2009.

SPECIAL CIRCUMSTANCES: SEVERE PREECLAMPSIA AT < 34 WEEKS

Laurence E. Shields, MD, Marian Regional Medical Center/Dignity Health

BACKGROUND:

Preeclampsia presenting at < 37 weeks:

The criteria for diagnosing mild and severe preeclampsia are the same regardless of gestational age (Table 1).¹ Preeclampsia is generally a disorder that manifests near term with an overall incidence of 5-6% of all pregnancies.¹ Preeclampsia is also seen earlier in gestation, especially when other co-morbidities exist (e.g., diabetes, chronic hypertension, systemic lupus erythematosus).^{2,3} Development of preeclampsia <37 weeks is seen in approximately 1.5% of births and about 50% of these cases represent severe disease.^{2,4,5} About 0.3% of pregnant women will develop severe preeclampsia at <34 weeks.² For California, with an estimated 500,000 births annually this equates to approximately 1,500 women per year that will develop severe disease prior to 34 weeks gestation. For patients with mild disease ≥ 37 weeks or severe disease ≥ 34 weeks, delivery is indicated as the balance of maternal risk and potential neonatal benefit favors delivery.⁶⁻⁸

Severe Preeclampsia presenting < 34 weeks:

Delivery has been traditionally recommended for patients with severe preeclampsia presenting at any gestational age. However, a number of observational studies and a limited number of randomized clinical trials suggest that under appropriate conditions, patients with severe disease at < 34 weeks gestation may be conservatively managed.^{4,6} These patients will be primarily those with blood pressure that can be controlled in a relatively short period of time, **without** manifestations of end-organ disease. Significant proteinuria, an indication of renal involvement, is the one exception to the recommendation for immediate delivery. Normal fetal testing is a prerequisite for conservative management.⁶⁻⁸ Severe IUGR with reassuring fetal monitoring is a clinical scenario in which conservative management is a reasonable option. These patients should be hospitalized and their status must be continuously assessed for disease progression. Delivery is indicated if there is evidence of fetal or maternal deterioration.

Treatment of maternal blood pressure should be maintained at a level that reduces the risk of maternal cerebral vascular accidents (diastolic BP < 105-110 mmHg and systolic BP <160 mmHg).^{9,10} The level of maternal blood pressure control and recommended medications are outlined in the Medications document. Initial treatment should include treatment with magnesium sulfate 4-6 gram over 15-20 minutes then 1-2 gm per hour (see section on magnesium sulfate). Betamethasone for induction of fetal lung maturity and decreasing the risk of neonatal complications such as Respiratory Distress Syndrome (RDS), Intraventricular hemorrhage (IVH), and necrotizing enterocolitis (NEC), should also be given at the time of admission.

Maternal monitoring should be carried out frequently (Table 2) in a facility that has the capacity to take care of both the mother and the infant, with equipment, personnel and neonatal care facilities that can adequately care for infants delivered at a gestational age of ≤ 34 weeks. Complications common to this group of women include seizure, stroke, Posterior Reversible Encephalopathy Syndrome (PRES), pulmonary edema, Adult Respiratory Distress Syndrome (ARDS), placental abruption, liver rupture, obstetrical hemorrhage, Disseminated Intravascular Coagulation (DIC), and acute renal failure.² Assessment of maternal central nervous system (CNS): headache, visual changes or change in mental status) and gastrointestinal (GI): epigastric, Right Upper Quadrant, [RUQ], pain or nausea and vomiting) symptoms should be carried out with each maternal vital sign assessment, no longer appropriate and delivery should be instituted.

Most expert opinions^{6,7,9} and other international organizations^{11,12} state that delaying delivery is not recommended in the presence of the following: 1) Eclampsia; 2) HELLP (Hemolysis, Elevated Liver Enzymes, Low Platelet Syndrome); 3) Pulmonary edema; 4) Severe thrombocytopenia; or 5) Coagulopathy (Table 1). *Any patient who does not meet the criteria for continuation of pregnancy should be delivered by either induction of labor or cesarean section.* Determining the route of delivery should take into consideration the likelihood of success based on cervical status, gestational age, status of the fetus, and the severity of the disease.

Each obstetrical unit should develop a policy that delineates the conditions under which mothers and neonates can be effectively treated at that institution. A decision checklist specific to the level of care that can be provided should be used to assist physicians with decisions related to transfer. If the mother or the neonate can not be adequately cared for at the center where she presents, strong consideration should be given to transport to a center that can provide a higher level of care.

A multidisciplinary working group that includes: Obstetrician, Obstetrical nursing, Anesthesia, Maternal Fetal Medicine (MFM), (if available), Intensivist and Intensive Care Unit (ICU) nursing as needed should coordinate the patient's care. The patient's care plan should address the specific problems presented by the patient's specific manifestations of preeclampsia, as well as anticipated complications. It is likely that the obstetrician, obstetrical nurse or MFM specialist will need to take the lead in this arena. The pediatric/neonatal team should be involved to provide patient consultation, education, and anticipatory outcome guidance based on gestational age. The patient also needs to be aware, that continuation of the pregnancy is being attempted with the goal of improving fetal outcome. However, the patient also needs to understand that there is a risk, albeit low, of fetal demise and/or significant maternal morbidity.

Table 1. Severe Preeclampsia and Management Options for Delayed Delivery.^{6,7}

Criteria	Definition/Significance	Attempt to Delay Delivery
Persistent headache/blurred vision or scotomata*/mental status changes**	Suggest central nervous system dysfunction	No
Persistent epigastric pain or right upper quadrant pain	Suggest liver capsule distension or rupture	No
Eclampsia	Generalized tonic clonic seizure	No
Pulmonary edema and/or hypoxia (O2 saturation < 95%)	Excessive fluid accumulation in the lungs	No
Oliguria/Renal failure	Urine output of <500/24 hours or Creatinine >1.2 (unless chronic renal disease)	No
Hepatocellular injury	Serum transaminases >2x normal	No
Blood Pressure	>160/110 mmHg BP criteria for Severe Preeclampsia	Yes, if responds to treatment
Proteinuria	> 5 grams/24 hours	Yes
Intrauterine growth restriction	With reassuring fetal testing***	Yes

*Patients with eclampsia and visual disturbances should be evaluated in consultation with critical care medicine/neurology for the presence of Posterior Reversible Encephalopathy Syndrome (PRES).

** Mental status changes in the presence of severe thrombocytopenia should be evaluated in consultation with hematology for Thrombotic Thrombocytopenic Purpura (TTP) and consideration for treatment or transfer to a center with treatment capacity should be given.

*** Nonstress test, Contraction Stress Test (CST), or Biophysical Profile (BPP) carried out twice daily or every shift.

Table 2. Daily Assessment for Delivery versus Continuing Pregnancy
IMMEDIATE DELIVERY (post maternal stabilization) Indicated for:

Criteria	Definition/Significance
Maternal headache/blurred vision or Scotomata*/mental status/changes**	Suggest central nervous system dysfunction
Persistent epigastric pain or right upper quadrant pain	Suggest liver capsule distension or rupture
Eclampsia	Generalized tonic clonic seizure
Pulmonary edema or hypoxia	Excessive fluid accumulation in the lungs
O2 saturation <95%	
Oliguria/renal failure	Urine output of <500/24 hours or Creatinine > 1.2
Hepatocellular injury	Serum transaminases >2x normal

If laboratory values were normal at admission and remain normal for two consecutive days and blood pressure is stable (i.e. not requiring additional medication) then every other to every third day laboratory monitoring may be used.

Hemolysis, Elevated Liver Enzymes, Low Platelets (HELLP) Syndrome:

HELLP syndrome is a variant of severe preeclampsia characterized by red blood hemolysis, thrombocytopenia, and abnormal elevations in liver transaminases.¹³ The diagnostic criteria are listed in Table 3 below. Three classes of HELLP are characterized by severity of laboratory abnormalities and risk for significant adverse perinatal outcome based on the patient’s platelet count.¹⁴ The most severe manifestation (Class I) has platelet counts $\leq 50,000$ cells/ μL ; Class II has platelet counts of $>50,000$ and $\leq 100,000$ cells/ μL , and in Class III, there is mild thrombocytopenia with a platelet nadir between $>100,000$ and $\leq 150,000$ cells/ μL . The severity of maternal, fetal and neonatal morbidity is correlated with the severity of the disease.¹⁴ Approximately 10-15% of patients with classic HELLP syndrome will not have elevated blood pressures ($\text{BP} \geq 140/90$ mmHg)¹⁵ and like other forms of severe preeclampsia, proteinuria is absent in 15-25% of patients.¹⁴ The presence of subjective symptoms is seen in 64-84% of patients with Class III and Class I HELLP respectively.¹⁴ Thus, the presence of proteinuria or elevated blood pressure is not essential for the diagnosis of HELLP syndrome and in those patients without classic features, the presence of subjective symptoms (i.e. headache, epigastric pain, nausea and vomiting, or visual disturbances) should prompt further evaluation to rule out progression of disease requiring delivery.

Table 3: Classes of HELLP Syndrome

Class	Description
I	Platelet counts <50,000 cells/ μ L
II	Platelet counts >50,000 and <100,000 cells/ μ L
III	Mild thrombocytopenia Platelet nadir between >100,000 and <150,000 cells/ μ L

The majority of patients with HELLP syndrome will have elevated blood pressure spanning the range from mild to severe. The combination of severely elevated blood pressure with thrombocytopenia and abnormal coagulation parameters place the patient at increased risk for cerebral vascular accidents or other hemorrhagic complications. The frequency of seizures/eclampsia ranges from 5-12% in severe preeclampsia and HELLP Syndrome of any degree. (Class I, II, and III).¹⁴

Maternal and neonatal morbidity is significantly increased in pregnancies complicated by HELLP syndrome. The rate of preterm birth is high (70%) with 15% of deliveries occurring prior to 28 weeks.¹⁵ The risk of maternal death has been estimated at 1%, and the frequency of other severe morbidities is also high including Disseminated Intravascular Coagulation (DIC): 15-30%; Pulmonary Edema: 8%; Acute Renal Failure: 3% and; Adult Respiratory Distress Syndrome (ARDS) and Stroke: both 1%. Initiation of corticosteroid therapy to decrease maternal morbidity for Class I and II HELLP Syndrome should be considered.¹⁶

Table 4. Diagnostic Criteria for HELLP Syndrome.¹⁷

Hemolysis	Elevated LDH (>600 IU/L)* , Microangiopathic hemolytic anemia on peripheral blood smear, low haptoglobin (<25 mg/dL), elevated indirect bilirubin
Thrombocytopenia	Platelet count \leq 100,000 cells/ μ L
Elevated Transaminases	Serum AST \geq 70 IU/L or twice baseline values

*Elevated LDH >600IU/L is currently the most readily available and accurate laboratory indicator of hemolysis.

Diagnosis of HELLP syndrome can be challenging, as the differential diagnosis includes Thrombotic Thrombocytopenic Purpura (TTP), Hemolytic Uremic Syndrome (HUS), and Acute Fatty Liver Disease of Pregnancy (AFLP). TTP-HUS should be considered in all pregnant women with severe thrombocytopenia, severe anemia, and elevated LDH (lactic dehydrogenase) levels with minimal elevation of AST.¹⁸ A history of proteinuria and hypertension prior to onset of hemolysis, liver abnormalities, and thrombocytopenia favor the diagnosis of preeclampsia, while high LDH levels with only modest elevation of AST favors TTP (Table 4). The distinction between TTP or HUS and severe

preeclampsia or HELLP is important for therapeutic and prognostic reasons, as TTP would normally be treated with plasmapheresis.

Table 5. Differentiation between Preeclampsia, HELLP Syndrome, Acute Fatty Liver Disease of Pregnancy (AFLD), Thrombotic Thrombocytopenia Purpura (TTP), Hemolytic Uremia Syndrome (HUS). **

	Plts	LFTs	Bili	Cr	LDH	Glu	DIC	CNS
Preeclampsia	±	±	±	±	±	→	±	±
HELLP	↓/↓↓	↑↑	↑	±	↑	→	±	±
AFLD	↓↓	↑↑	↑↑↑	↑	↑	↓↓↓	↑↑↑	±
TTP	↓↓↓	↑	↑	↑	↑↑	→	±	++
HUS	↓	↑↑	↑↑	↑↑↑	↑	→	±	±

AFLD: Acute Fatty Liver Disease of Pregnancy; TTP: Thrombotic Thrombocytopenic Purpura; HUS: Hemolytic Uremia Syndrome; Plts: platelet count; LFTs: liver transaminases; Bili: total bilirubin level; LDH: Lactate Dehydrogenase; Glu: glucose; DIC: Disseminated Intravascular Coagulation; CNS: Central Nervous System symptoms (confusion, visual changes, headache)

**Arrows represent relative changes: one arrow equals some increase; two arrows indicate moderate increase, and three arrows equal very high increase

Hypertension:

Other health systems have defined the degree of hypertension associated with pregnancy into three categories (mild, moderate, and severe, see Table 5 below).¹⁹ The rationale for this expanded categorization is to recognize that many patients in the moderate hypertension category are at higher risk for poor obstetrical outcome and likely merit closer observation.

Table 6. Preeclampsia Diagnostic Criteria for Three Blood Pressure Categories (mmHg)

SOGC*, NICE**	Mild	Moderate	Severe
Systolic	140-149	150-159	>160
Diastolic	90-99	100-109	>110

*SOGC (Society of Obstetricians and Gynaecologists of Canada)

**NICE (National Institute for Health and Clinical Excellence)

Table 7: Preeclampsia Diagnostic Criteria for Two Blood Pressure Categories (mmHg)

ACOG*	Mild	Severe
Systolic	140-159	>160
Diastolic	90-109	>110

*ACOG (American College of Obstetrics and Gynecology)

Atypical Preeclampsia:

The recognition that the pathophysiology of preeclampsia is highly variable has led to the realization that the disease can present with single- or multi-organ dysfunction.¹⁸ Cases that present as “atypical” are those that manifest at <20 weeks gestation, more than 48 hours after birth or with any of the diagnostic criteria for severe disease in the absence of proteinuria or elevated blood pressure. Severe nausea and vomiting in the late 2nd or early 3rd trimester should raise the index of suspicion for preeclampsia. Development of preeclampsia in the presence of gestational hypertension is inversely related to the time of diagnosis of gestational hypertension.²⁰ Furthermore, proteinuria, in the absence of hypertension, may be the first manifestation of disease in the sequence leading to preeclampsia.²¹ These women merit close antenatal follow-up (1-2 times per week) with laboratory assessment.¹⁸ Women presenting after delivery with gestational hypertension or isolated proteinuria and who have laboratory or subjective symptoms of severe preeclampsia should be treated with magnesium sulfate.

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. Patients with diagnostic criteria for mild preeclampsia at ≥ 37 weeks or severe preeclampsia at ≥ 34 weeks gestation should be delivered.
2. Patients with diagnostic criteria for severe preeclampsia at <34 weeks gestation should be delivered if criteria outlined in Table 1 (above) are not met.
3. Patients with severe preeclampsia remote from term (<34 weeks) should be managed at or transported to a center with experience and expertise in management of these patients as well as their potential complications.
4. Blood pressure should be controlled to a level that is between 140-160 mmHg systolic and 90-105 mmHg diastolic (See medication section)
5. Magnesium sulfate should be used in all patients with HELLP syndrome, severe preeclampsia, or unstable patients whose disease is evolving.
6. Mode of delivery should be based on usual obstetrical indications.
7. Corticosteroids may be considered in cases of HELLP syndrome for decreasing maternal morbidity (see Medication, [pg. 48,] and Antihypertensive, [pg 45] chapters).²²
8. Patients with moderate hypertension (150-159 mmHg systolic and 100-109 mmHg diastolic) with and without proteinuria should be monitored with a heightened level of

supervision, including the frequency of blood pressure measurements, laboratory studies and symptom assessment. Antihypertensive therapy should be considered in the group of patient that have blood pressures that are >155 systolic and 105 diastolic.¹⁰

9. Patients with elevated blood pressure not having proteinuria but who have other diagnostic criteria of severe preeclampsia should be treated as if they have severe disease.

EVIDENCE GRADING

Level of Evidence: II-1A, III-1A

REFERENCES

1. American College of Obstetricians and Gynecologists. Diagnosis and management of preeclampsia and eclampsia; ACOG Practice Bulletin No. 33. *Obstet & Gynecol.* January 2002, Reaffirmed 2010;99:159-167.
2. Gupta L, Gaston L, Chauhan S. Detection of fetal growth restriction with preterm severe preeclampsia: experience at two tertiary centers. *Am J Perinatol.* 2008;25:247-249.
3. Hutcheon J, Lisonkova S, Joseph K. Epidemiology of pre-eclampsia and the other hypertensive disorders of pregnancy. *Best Pract Res Clin Obstet Gynaecol.* 2011;25:391-403.
4. Sibai BM. Evaluation and management of severe preeclampsia before 34 weeks' gestation. *Am J Obstet Gynecol.* Sep 2011;205(3):191-198.
5. Catov J, Ness R, Kip K, Olsen J. Risk of early or severe pre-eclampsia related to pre-existing conditions. *Int J Epidemiol.* 2007;36:412-419.
6. Norwitz E, Funai E. Expectant management of severe preeclampsia remote from term: hope for the best, but expect the worst. *Am J Obstet Gynecol.* 2008;199:209-212.
7. Sibai B. Management of late preterm and early-term pregnancies complicated by mild gestational hypertension/pre-eclampsia. *Semin Perinatol.* 2011;35:292-296.
8. Haddad B, Deis S, Goffinet F, Paniel BJ, Cabrol D, Sibai B. Maternal and perinatal outcomes during expectant management of 239 severe preeclamptic women between 24 and 33 weeks' gestation. *Am J Obstet Gynecol.* 2004;190:1590-1595; discussion 1595-1597.
9. ACOG. Committee Opinion no. 514: Emergent therapy for acute-onset, severe hypertension with preeclampsia or eclampsia. *Obstet Gynecol.* 2011;118:1465-1468.
10. Martin J, Thigpen B, Moore R, Rose C, Cushman J, May W. Stroke and severe preeclampsia and eclampsia: a paradigm shift focusing on systolic blood pressure. *Obstet & Gynecol.* 2005;105(2):246-254.

11. Visintin C, Mugglestone M, Almerie MQ, Nherera L, James DC, Walkinshaw S. Management of hypertensive disorders during pregnancy: summary of NICE guidance. *BMJ*. 2010;25(341(25)):c2207.
12. Chaillet N, Dubv/© E, Dugas M, et al. Identifying barriers and facilitators towards implementing guidelines to reduce caesarean section rates in Quebec. *Bull World Health Organ*. 2007;85(10):791-797.
13. Weinstein L. Syndrome of hemolysis, elevated liver enzymes, and low platelet count: a severe consequence of hypertension in pregnancy. *Am J Obstet Gynecol*. 1982;142:159-167.
14. Martin J, Rinehart B, May W, Magann E, Terrone D, Blake P. The spectrum of severe preeclampsia: comparative analysis by HELLP (hemolysis, elevated liver enzyme levels, and low platelet count) syndrome classification. *Am J Obstet Gynecol*. 1999;180:1373-1384.
15. Sibai B. Diagnosis, controversies and management of HELLP syndrome. *Obstetrics & Gynecology*. 2004;103:981-991.
16. Martin J, Owens M, Keiser S, et al. Standardized Mississippi Protocol treatment of 190 patients with HELLP syndrome: slowing disease progression and preventing new major maternal morbidity. *Hypertens Pregnancy*. 2012;31(1):79-90.
17. Sibai B, Ramadan M, Usta I, Salama M, Mercer B, Friedman S. Maternal morbidity and mortality in 442 pregnancies with hemolysis, elevated liver enzymes, and low platelets (HELLP Syndrome). *American Journal of Obstetrics and Gynecology*. 1993;169:1000-1006.
18. Sibai BM, Stella CL. Diagnosis and management of atypical preeclampsia-eclampsia. *American Journal of Obstetrics and Gynecology*. May 2009;200(5):481 e481-487.
19. National Collaborating Centre for Women's and Children's Health (UK). Hypertension in pregnancy the management of hypertensive disorders during pregnancy. *NICE Clinical Guidelines, No. 107*: RCOG Press; 2010.
20. Barton J, O'Brien J, Bergauer N, Jacques D, Sibai B. Mild gestational hypertension remote from term: progression and outcome. *Am J Obstet Gynecol*. 2001;184:979-983.
21. Airoldi J, Weinstein L. Clinical signifiacne of proteinuria in pregnancy. *Obstet Gynecol*. 2007;Survey 62:11-24.
22. Martin J, Brewer J, Wallace K, et al. HELLP Syndrome and composite major maternal morbidity: importance of Mississippi classification system. *Maternal Fetal Neonatal Med*. 2013;Epub ahead of print DOI: 0.3109/14767058.2013.773308(Mar 6).

CONSULTATION TRIGGERS IN SEVERE PREECLAMPSIA FOR ALL OBSTETRIC UNITS

Mark Zakowski, MD, Cedars Sinai Medical Center

BACKGROUND

Patients with preeclampsia are at risk for numerous adverse outcomes. The labor and delivery team of obstetricians, nurses and anesthesiologists are the first responders, but will require consultation with other specialties in a number of clinical circumstances. The following are guidelines for engaging additional practitioners in providing added clinical depth for patient care.

Table 1: Trigger Criteria for Consultations

Pulmonary/Fluids	Cardiac	Neurologic	Hematologic
<ul style="list-style-type: none"> • Pulmonary edema • Fluid overload, leaky membrane, low Colloid Oncotic Pressure • Not responding to one dose of diuretic • Shortness of breath– DDx includes r/o pulmonary embolism (spiral CT scan preferred) 	<ul style="list-style-type: none"> • Cardiac pump failure –(DDx) includes peripartum cardiomyopathy, preeclampsia induced – need echo. • Arrhythmia (e.g. SVT, atrial fibrillation) • Difficulty breathing, (might need intubation: DDx: pulmonary edema, stridor from swelling fluids/allergic, asthmatic not responsive to initial medications, magnesium toxicity, occult Mitral Stenosis for new onset asthma in labor • Hypoxia, any cause (decreased PO₂) – (e.g. oxygen saturation <95% on oxygen). Trauma history (possible pneumothorax – chest tube required) • Intrinsic – cardiac pump failure, leaky membrane, COP low, bronchospasm, Extrinsic – PTX, ETT kink, FB in airway, Swelling/stridor – fluid/preeclampsia progression labor, allergic reaction 	<ul style="list-style-type: none"> • Repeated seizures, unresponsive to initial therapy (DDx includes SAH/intracranial hemorrhage – CT required) • Altered mental status (DDx – metabolic, toxic, etc.) • Acute stroke/neurologic changes (r/o intracranial bleed) • Cortical vein thrombosis 	<ul style="list-style-type: none"> • DIC • HELLP syndrome (e.g. platelets <50,000) • Coagulopathy, any cause • Massive transfusion/OB hemorrhage • On anticoagulants (e.g., LMWH) – timing dosing, when to hold, when to restart

DDx: Differential Diagnosis; r/o: Rule Out; CT: computed tomography; SVT: Supraventricular Tachycardia; COP: colloid osmotic pressure; PTX; ETT; FB; DIC: Disseminated Intravascular Coagulopathy; HELLP: Hemolysis, Elevated Liver Enzymes, Low Platelet; LMWH: low molecular weight heparin

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. Consultation of maternal fetal medicine, anesthesia, cardiology, hematology, and/or neurology or any other sub-specialties should be strongly considered if the staff feel uncomfortable with the medical situation or if any hematologic, cardiac, pulmonary, or persistent neurologic symptoms are present.
2. Request consultations when the patient needs a higher level of care than usually provided by regular L and D staff, or the staff feel uncomfortable with the medical situation. Often the first consults are with the MFM and/or Anesthesiologist covering OB.^{1 2}
3. Consultations should also be considered in the following situations:
 - There is clinical disagreement among team members about the severity of the woman's condition
 - Hypertension is resistant to standard treatment (e.g., SBP>160mmHg, DBP>105 mmHg), need 3rd line drug (i.e., after labetalol, hydralazine per CMQCC/ACOG protocols)
 - Persistent low BP (e.g., SBP <90mmHg) unresponsive to fluid bolus(es) of 500 ml
 - Crystalloid and/or short acting vasopressors (e.g. ephedrine due to neuroaxis blockade)
 - Persistent oliguria (e.g., <30 cc per hr) after fluid challenge (See Fluid Management section, pg. 70)
 - Suspected amniotic fluid or pulmonary embolism, or;
 - Hemorrhage with disseminated intravascular coagulation (DIC).

EVIDENCE GRADING

Level of Evidence: C

REFERENCES

1. Kodali B, Chandrasekhar S, Bulich L, Topulos G, Datta S. Airway changes during labor and delivery. *Anesthesiology*. 2008;108:357-362.
2. Isono S. Mallampati classification, an estimate of upper airway anatomical balance, can change rapidly during labor. *Anesthesiology*. 2008;108:347-349.

POSTERIOR REVERSIBLE ENCEPHALOPATHY SYNDROME (PRES)

Racine Edwards-Silva, MD, Olive View-UCLA Medical Center

BACKGROUND

Posterior reversible encephalopathy syndrome (PRES) is a transient clinical neuroradiological entity characterized by clinical signs and symptoms including hypertension, generalized seizure activity, altered mental status, headache and vision changes; along with PRES findings characteristic on head computed tomography (CT) or magnetic resonance imaging (MRI) scan.¹ Many causes of PRES have been reported in the literature including hypertensive encephalopathy, preeclampsia, eclampsia, renal failure, immunosuppressants, thrombotic thrombocytopenic purpura, systemic lupus erythematosus (SLE), and acute intermittent porphyria.² The nomenclature for this syndrome has undergone several changes with one radiologic journal describing this entity as Eclamptic Encephalopathy.³ In the early postpartum period PRES is most often seen in association with severe elevated BPs, with eclampsia, or in the Emergency Department with a patient who presents in the late postpartum period with hypertension and seizures.


Preeclampsia and eclampsia are probably the most common causes of PRES. The true incidence of PRES complicating preeclampsia and eclampsia is unknown because neuroradiographic imaging is not routinely performed.

Symptoms: Neurologic symptoms commonly precede eclampsia. Headache and visual disturbance are the most common prodromal symptoms. These neurologic symptoms reflect the development of cerebral edema and vasospasm of cerebral and retinal vessels. Premonitory symptoms may provide an early warning of imminent eclampsia. Other common warning symptoms include nausea, vomiting, or epigastric/abdominal pain. These gastrointestinal symptoms are believed to reflect hepatocellular involvement from periportal or parenchymal necrosis, liver capsule stretching and/or hemorrhage.⁴


Treatment: PRES is usually reversible with prompt diagnosis and treatment. Early recognition and effective treatment of blood pressure in patients with PRES in the acute setting along with seizure prophylaxis decreases long-term sequelae of this condition. The antepartum patient should be stabilized and then delivered.

However, in rare cases the reversible vasogenic edema associated with PRES can progress to irreversible ischemic damage, cerebral infarction or even death. Treatment is the same as for severe preeclampsia, and neurological consultation is recommended. The underlying pathophysiology has been attributed to failure of cerebral auto-regulation and endothelial dysfunction. This impairment of cerebral auto-regulation leads to disruption of the blood-brain barrier in the posterior circulation with resultant extravasation of fluids and protein across the altered blood-brain barrier.⁵ This process

causes the characteristic lesions seen in the occipital and posterior parietal lobes on neuroradiologic imaging (Photos A, B, C). These radiographic cerebral abnormalities appear as intense signals on T2-weighted MRI scans and as low-density areas on CT scans. (Photos are of same patient taken at the same time, different views.)



A



Photo

Photo B




Photo C

Photos highlight involvement of the occipital region that explains the visual changes seen in preeclampsia/PRES. Photos used with kind permission of Thomas Archer, MD, University of California, San Diego, 2013

Diagnosis: MRI is considered the most appropriate tool as an adjunct in the clinical setting to diagnosing PRES and demonstrating the characteristic brain lesions. MRI

imaging is superior to CT imaging in patients with PRES or Eclamptic Encephalopathy. The hallmark feature is bilateral symmetrical vasogenic edema in the territories of the posterior cerebral circulation white matter (occipital and posterior parietal lobes). The posterior cerebral white matter edema is most evident on T2-weighted MRI images with fluid-attenuated inversion recovery (FLAIR).⁶ The predominance of occipital lesions corresponds well to the neurologic manifestation of temporary cortical blindness. Neuroradiologic imaging should be strongly considered in the postpartum period with a patient who presents with headache, hypertension, seizures, or atypical neurologic symptoms such as visible changes or inability to see (blindness), in order to rule out other differential diagnoses such as a mass lesion or cerebral venous thrombosis.

KEY LEARNING POINTS

1. Upon initial discharge from the hospital, inform the recently delivered patient that there is still a measurable risk of Preeclampsia/Eclampsia in the postpartum period for up to six (6) weeks after delivery, even with an uneventful pregnancy and delivery. Although delayed postpartum preeclampsia is usually seen within the first two weeks postpartum, some patients can present up to 4-6 weeks after delivery. This risk may be increased in patients who have experienced preeclampsia prior to delivery. Therefore, delayed preeclampsia should be considered in any patient within the normal six-week postpartum period.
2. It is important for Emergency Department physicians/staff to have a high index of suspicion for postpartum Preeclampsia/Eclampsia and to have early involvement of obstetric staff in the care of these patients.
3. Upon presentation to the ED, a female patient should be queried as to whether she is currently pregnant or has recently been pregnant.
4. Antihypertensive medications should be implemented for systolic BP >160 mmHg and/or if diastolic BP >105-110 mmHg that persists for 15 minutes or greater and should be considered in the group of patients that have blood pressures that are >155 systolic and 105 diastolic.⁷
5. Treatment guidelines for severe hypertension are identical to those for severe preeclampsia/eclampsia.

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. Initiate established treatment algorithms for severe preeclampsia and eclampsia for control of hypertension and seizures.
2. First-line antihypertensive medication is intravenous Labetalol or Hydralazine. (See Antihypertensive Agent chapter. (pg. 45)
3. Magnesium sulfate is the treatment of choice for controlling seizures in eclampsia and for prevention of recurrent seizures. This should be maintained for at least 24 hours after the last seizure.(See Seizure Medication chapter: Magnesium Sulfate, pg. 48)
4. If patient is pregnant, maternal and fetal monitoring is advised on L&D as fetal status is secondary to maternal stabilization.
5. In these complex clinical cases, a multidisciplinary team of ED, OB, Anesthesia, and possibly Neurology or Critical Care is necessary.
6. Do not delay treatment to perform neuroradiologic imaging.
7. Neuroradiologic imaging is strongly advised in the postpartum period due to the numerous differential diagnoses and to exclude other intracranial pathology.

EVIDENCE GRADING

Level of Evidence: III

REFERENCES

1. Powell E, Goldman M. Posterior reversible encephalopathy syndrome (PRES) in a thirty-six week gestation eclamptic. *Emerg Med.* 2007;Nov(33(4)):377-379.
2. Cho H, Lee H. Posterior reversible encephalopathy syndrome in early postpartum woman. *Hong Kong Journal of Emerg Med.* 2012;19(1):58-61.
3. Zak I, Dulai H, Kish K. Imaging of Neurologic Disorders Associated with Pregnancy and the Postpartum Period. *RadioGraphics.* 2007;27(1):85-108.
4. Cooray S, Edmonds S, Tong S, Samarasekera S, Whitehead C. Characterization of Symptoms Immediately Preceding Eclampsia. *Obstet Gynecol.* 2011;118(5):995-999.
5. Nuwer J, Eshaghian S. Late Postpartum Eclampsia with Posterior Reversible Encephalopathy Syndrome. *Hospital Physician.* 2007:45-49.

6. Long T, Hein B, Brown M, Rydberg C, Wass C. Posterior reversible encephalopathy syndrome during pregnancy: seizures in a previously healthy parturient. *J Clin Anes.* 2007;19:145-148.
7. Martin J, Thigpen B, Moore R, Rose C, Cushman J, May W. Stroke and severe preeclampsia and eclampsia: a paradigm shift focusing on systolic blood pressure. *Obstet & Gynecol.* 2005;105(2):246-254.

EMERGENCY DEPARTMENT and NON-OBSTETRICAL VISITS

EMERGENCY DEPARTMENT RECOGNITION and TREATMENT: FOCUS ON DELAYED POSTPARTUM PREECLAMPSIA and ECLAMPSIA

Mark Meyer, MD, Southern California Kaiser Permanente Medical Group

BACKGROUND:

Hypertensive disorders including preeclampsia and eclampsia are one of the leading causes of maternal morbidity and mortality. While there has been an overall decrease in the frequency of eclampsia, the frequency of postpartum and delayed eclampsia has increased¹ making it more common for patients to present to the Emergency Department (ED) with symptoms. Postpartum or delayed preeclampsia/eclampsia is frequently associated with PRES (Posterior Reversible Encephalopathy Syndrome). Although obstetric consultation is warranted in every case of preeclampsia, emergency physicians should be knowledgeable of and comfortable with the initial management. Since many of these patients will present to an ED, education of ED personnel and application of diagnosis and treatment protocols are important steps in reducing morbidity and mortality associated with postpartum preeclampsia and eclampsia. Emergency physicians should have a higher index of suspicion in order to improve the recognition and treatment of postpartum preeclampsia and eclampsia. This may require gathering historical information regarding a recent pregnancy from family members; it is important to remember that:

1. Up to 26% of eclamptic seizures occur beyond 48 hours and as late as four to six (6) weeks after delivery.^{1,2} However, most of these cases occur in the first seven (7) days after delivery.³
2. As many as 78% of these patients have no previous diagnosis of hypertensive disease with the antecedent pregnancy.^{2,3}
3. If medical records are not immediately available, treating personnel may have no knowledge that the patient has recently delivered, resulting in a decreased index of suspicion.⁴
4. While the clinical presentation of delayed postpartum preeclampsia may be atypical, the most common complaint is headache in up to 69% of patients.³ Headache in a recently pregnant patient will likely be isolated but should prompt an investigation into the possibility of delayed postpartum preeclampsia.

Seizures in the first and early second trimester (<20 weeks) or well into the postpartum period probably are due to Central Nervous System (CNS) pathology and warrant full evaluation, including computed tomography (CT) scanning of the head, lumbar puncture (if clinical evidence of meningitis or concern for hemorrhage exists), determination of electrolyte levels and urine or serum toxicologic screening. Do not overlook other

neurologic causes of seizure, particularly if the seizure occurs more than 24-48 hours after delivery.

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. ED triage protocols must identify patients who are currently pregnant or have delivered in the previous six (6) weeks. If the patient's medical records are not available, then simple questioning of the patient, family, Emergency Medical Services (EMS), etc., may provide this information. This information must then be clearly communicated to the treatment team.
2. ED personnel should be familiar with the risk factors and characteristics of delayed postpartum preeclampsia and eclampsia.
3. Do not overlook other neurologic causes of seizure, particularly if the seizure occurs more than 48 hours after delivery.
4. Implementation of the CMQCC (see Appendix F) protocol for diagnosis and treatment of preeclampsia and eclampsia in the Emergency Department. This can be reinforced through the use of educational tools in other sections of this toolkit and with the use of drills and simulations.


EVIDENCE GRADING

Level of Evidence: C

REFERENCES

1. Chames M, Livingston J, Ivester T, Barton J, Sibai B. Late postpartum eclampsia: a preventable disease? *Am J Obstet Gynecol.* 2002;186:1174-1177.
2. Matthys L, Coppage K, Lambers D, Barton J, Sibai B. Delayed postpartum preeclampsia: an experience of 151 cases. *Am J Obstet Gynecol.* 2004;190:1464-1466.
3. Al-Safi Z, Imudia A, Filetti L, Hobson D, et al. Delayed postpartum preeclampsia and eclampsia: demographics, clinical course, and complications. *Obstet Gynecol.* 2011;118(5):1102-1107.
4. The Joint Commission. Preventing Maternal Death. Sentinel Event Alert. Issue 44. 2010;
http://www.jointcommission.org/sentinal_event_alert_issue_44_preventing_maternal_death. Accessed January 26, 2010.

Part 1 of 2: Diagnosis - Evaluation and Treatment of Antepartum and Postpartum Preeclampsia and Eclampsia in the Emergency Department



Part 2 of 2: Treatment - Evaluation and Treatment of Antepartum and Postpartum Preeclampsia and Eclampsia in the Emergency Department

Evaluation and Treatment of Antepartum and Postpartum Preeclampsia and Eclampsia in the Emergency Department

TREATMENT

1st Line Anti-Hypertensive Treatment: Labetalol & Hydralazine*
Target BP: 140-160/90-100/90 (BP<140/90 = decreased fetal perfusion)
See CMQCC Preeclampsia Toolkit for "Antihypertensives in Preeclampsia" for 2nd line therapy

Magnesium

LABELTALOL as Primary Anti-Hypertensive

- Administer Labetalol 20 mg IV
- Repeat BP in 10 min
 - If BP threshold is still exceeded, administer labetalol
 - If SBP<160 and DBP<100, continue to monitor closely
- Repeat BP in 10 min
 - If BP threshold is still exceeded, administer labetalol 80 mg IV
 - If SBP<160 and DBP<100, continue to monitor BP closely
- Repeat BP in 10 min
 - If BP threshold is still exceeded, administer hydralazine 10 mg IV
 - If SBP<160 and DBP<100, continue to monitor closely
- Repeat BP in 20 min; if BP threshold is still exceeded, obtain emergent consultation from maternal-fetal medicine, internal medicine, anesthesiology, or critical care
- Once target BP achieved, monitor BP q10 min for 1 hour, q 15 min for 2nd hour

HYDRALAZINE as Primary Anti-Hypertensive

- Administer hydralazine 5 or 10 mg IV
- Repeat BP in 20 min
 - If BP threshold is still exceeded, administer hydralazine 10 mg IV
 - If SBP<160 and DBP<100, continue to monitor closely
- Repeat BP in 20 min
 - If BP threshold is still exceeded, administer labetalol 20 mg IV
 - If SBP<160 and DBP<100, continue to monitor BP closely
- Repeat BP in 10 min
 - If BP threshold is still exceeded, administer labetalol 40 mg IV and obtain emergent consultation from maternal-fetal medicine, internal medicine, anesthesiology, or critical care
 - If SBP<160 and DBP<100, continue to monitor closely
- Once target BP achieved, monitor BP q10 min for 1 hour, q 15 min for 2nd hour

Initial Treatment

- Loading Dose: 4-6 gm over 15-20 min
- Maintenance 1-2 gm/hr
- Close observation for signs of toxicity
 - Disappearance of deep tendon reflexes
 - Decreased RR, shallow respirations, shortness of breath
 - Heart block, chest pain
 - Pulmonary edema

If Patient Seizes While on Magnesium:

- Secure airway and maintain oxygenation
- Give 2nd loading dose of 2 gm Magnesium over 5 min
- If patient seizes after 2nd magnesium bolus, consider the following:
 - Midazolam 1-2mg IV; may repeat in 5-10 min **OR**
 - Lorazepam 2 mg IV-may repeat **OR**
 - Diazepam 5-10 mg IV. May repeat q15 min to max of 30 mg
 - Phenytoin 1 g IV over 20 min

Seizures Resolve

- Maintain airway and oxygenation
- Monitor VS, cardiac rhythm/ECG for signs of medication toxicity
- Consider brain imaging for:
 - Head trauma
 - Focal Seizure
 - Focal neurologic findings
 - Other neurologic diagnosis is suspected

*Labetalol & Hydralazine recommendations based on 2011 ACOG Committee Opinion #514 and Practice Bulletin #33, Reaffirmed 2010

EDUCATION AND PATIENT INFORMATION

PRENATAL AND POSTPARTUM PATIENT COUNSELING OR EDUCATION*

Meredith Drews, Preeclampsia Foundation

Eleni Tsigas, PhD, Preeclampsia Foundation

*With acknowledgement of support from Whitney B.You, MD, MPH, LCDR, MC, USN, U.S. Naval Medical Center

BACKGROUND

Interventions for women with preeclampsia in the prenatal period include increased monitoring, magnesium sulfate, antihypertensive medications and corticosteroids for fetal lung maturation, if indicated. To maximally benefit from these resources, however, women must first seek medical care in a timely fashion.¹

Women are less likely to seek care if they have a poor understanding of the signs and symptoms of preeclampsia. Several recent studies emphasized the value of educating mothers and providers to report signs and symptoms of severe preeclampsia that commonly precede eclampsia, hypertensive encephalopathy, pulmonary edema or stroke.²⁻⁸ These recommendations are further supported by studies showing women who are diagnosed with preeclampsia, and receive timely and proper monitoring, have fewer adverse events than those with delayed diagnosis.⁴ This knowledge deficit appears modifiable, regardless of literacy level or initial understanding of preeclampsia, as pregnant women who had acknowledged receiving information about the disease, demonstrated greater preeclampsia-specific knowledge.⁹

Further, many clinicians and patients are unaware that preeclampsia can either occur or persist following delivery. It is also important to remember that the natural progression of postpartum hypertension includes an initial decrease in blood pressure (BP) within 48 hours, but BP rises again between 3-6 days postpartum.¹⁰ Preeclampsia may occur up to six (6) weeks postpartum.^{11,12} Postpartum hypertension/preeclampsia is either secondary to persistence or exacerbation of hypertension in women with previous gestational hypertension, preeclampsia, chronic hypertension or because of *de novo* (new onset) condition. In cases of late postpartum eclampsia, researchers found that nearly all of the patients had at least one prodromal symptom and half had more than one symptom that heralded the eclamptic seizure. Only 33% of women sought care for their symptoms, suggesting a need for proper patient education which may have led to better outcomes.³

KEY LEARNING POINTS

1. Many women have a limited understanding of preeclampsia, its signs and symptoms and its danger to both the mother and baby.¹³
2. Lack of understanding of preeclampsia and its prodromal symptoms is even more profound among women with low literacy levels.⁹
3. Health care providers may often overlook patients' complaints that in retrospect were predictors of increased risk or evidence of disease.
4. There is currently minimal education for a postpartum mother regarding preeclampsia at discharge from the hospital.
5. Many hospitals have discharge paperwork for obstetric patients that include warning signs that they should report to their doctors or that require immediate evaluation at a hospital. Symptoms of preeclampsia should be included in that list.
6. Many hospitals now also include videos on matters relating to new mothers that they may watch prior to discharge. These videos provide both a verbal and visual way to reinforce the warning signs of preeclampsia and what and when they need to communicate with their doctors.¹³
7. New mothers may often disregard symptoms since they may not know what they "should" be feeling postpartum. Family members are key partners in preventing maternal deaths by intervening when their spouse or partner complains of shortness of breath, relentless headache and other concerning symptoms.

RECOMMENDATIONS FOR QUALITY IMPROVEMENT:

1. A clear and simply written list of patient symptoms should be shared with expectant mothers and attending family members during prenatal visits and upon discharge from the hospital.¹⁴
2. A pictogram (Figure 1) showing the symptoms in visual format can be helpful to those women with language barriers or who may be struggling to understand the physician's instructions regarding preeclampsia.¹⁵
3. Physicians and nurses should ask open-ended questions to ensure that the patient understands what they have been told. For example, after going over list of symptoms say, "We've gone over a lot of information today. What would make you call or come in to the hospital?"¹⁶⁻²⁰

4. Hospitals with video education abilities should include a video on preeclampsia for patient education.
5. Women who have experienced preeclampsia prior to delivery or while in the hospital should see their OB within one week if they are on medication or two weeks if they are not on medications after discharge. A postpartum clinic visit should be established prior to discharge.
6. Women who have experienced preeclampsia prior to delivery or while in the hospital should be encouraged to monitor their blood pressure at home with instructions to call their physician if their pressures reach or exceed 140/90.
7. It is recommended that all patient discharge instructions (verbal and written) should include recognition of and response to preeclampsia symptoms and nurses should review these instructions with the patient and her family prior to discharge.

Figure 1. Preeclampsia Foundation Signs and Symptoms Information Sheet

Preeclampsia symptoms can be conveyed via a pictorial information sheet.⁹ (Note: This is available from the Preeclampsia Foundation for a modest shipping and handling fee. A Spanish version is also available.)

Ask Your Doctor or Midwife

Preeclampsia

What Is It?

Preeclampsia is a serious disease related to high blood pressure. It can happen to any pregnant woman.


Risks to You

- Seizures
- Stroke
- Organ damage
- Death


Risks to Your Baby

- Premature birth
- Death


Signs of Preeclampsia




Stomach pain




Headaches




Feeling nauseous; throwing up



Seeing spots



Swelling in your hands and face



Gaining more than 5 pounds in a week

What Should You Do?

Call your doctor right away. Finding preeclampsia early is important for you and your baby.

For more information go to www.preeclampsia.org

Copyright © 2010 Preeclampsia Foundation. All Rights Reserved.

REFERENCES


1. Goldenberg RL, McClure EM, Bhattacharya A, Groat TD, Stahl PJ. Women's perceptions regarding the safety of births at various gestational ages. *Obstet Gynecol.* Dec 2009;114(6):1254-1258.
2. Ogunyemi D, Benai J, Ukatu C. Is preeclampsia preventable? A case control review of consecutive cases from an urban underserved region. *South Med J.* 2004;97:440-445.
3. Chames M, Livingston J, Ivester T, Barton J, Sibai B. Late postpartum eclampsia: a preventable disease? *Am J Obstet Gynecol.* 2002;186:1174-1177.
4. Menzies J, Magee L, Li J, et al. Instituting surveillance guidelines and adverse outcomes in preeclampsia. *Obstet Gynecol.* 2007;110:121-127.
5. Sauve N, Powrie R, Larson L, et al. The impact of an educational pamphlet on knowledge and anxiety in women with preeclampsia. *Obstet Med.* 2008;1:11-17.
6. Matthys L, Coppage K, Lambers D, Barton J, Sibai B. Delayed postpartum preeclampsia: an experience of 151 cases. *Am J Obstet Gynecol.* 2004;190:1464-1466.
7. Filetti L, Imudia A, Al-Safi Z, Hobson D, Awonuga A, Bahado-Singh R. New onset delayed postpartum preeclampsia: different disorders? *J Matern Fetal Neonatal Med.* 2012;Jul(25(7)):957-960.
8. Al-Safi Z, Imudia A, Filetti L, Hobson D, et al. Delayed postpartum preeclampsia and eclampsia: demographics, clinical course, and complications. *Obstet Gynecol.* 2011;118(5):1102-1107.
9. You W, Wolf M, Bailey S, Grobman W. Improving patient understanding of preeclampsia: A randomized controlled trial. *Am J Obstet Gynecol.* 2012;206(5):431.e431-435.
10. Sibai BM. Etiology and management of postpartum hypertension-preeclampsia. *American Journal of Obstetrics and Gynecology.* Jun 2012;206(6):470-475.
11. Lubarsky S, Barton J, Friedman S, et al. Late postpartum eclampsia revisited. *Obstet Gynecol.* 1994;83:502-505.
12. Sibai B. Diagnosis, prevention, and management of eclampsia. *Obstet & Gynecol.* 2005;105(2):402-410.
13. Wilson E, Park D, Curtin L, et al. Media and memory: the efficacy of video and print materials for promoting patient education about asthma. *Patient Educ Couns.* 2010;80:393-398.
14. Freda M, Damus K, Merkatz I. Evaluation of the readability of ACOG patient education pamphlets. ACOG. *Obstet Gynecol.* 1999;May(93(5 Pt 1)):771-774.
15. MacGillivray I, McCaw-Binns A, Ashley D, Fedrick A, Golding J. Strategies to prevent eclampsia in a developing country: II. Use of a maternal pictorial card. *Int J Gynaecol Obstet.* 2004;87:295-300.

-
16. American Medical Association. Proceedings of 2005 White House Conference on Aging, Mini-Conference on Health Literacy and Health Disparities. 2005; <http://www.ama-assn.org/go/aging>. Accessed June 14, 2013.
 17. Institute of Medicine Health Literacy. *A Prescription to End Confusion*. Washington, DC: Institute of Medicine 2004.
 18. Ratzan S, Parker R. *National Library of Medicine Current Biographies in Medicine: Health Literacy*. Bethesda, MD: NCM;2000.
 19. The American Medical Association Foundation. Health Literacy and Patient Safety: Help Patients Understand. Reducing the risk by designing a safer, shame-free health care environment. 2007; http://www.ama-assn.org/resources/doc/ama-foundation/hl_monograph.pdf. Accessed December 7, 2011, 2007.
 20. Doak C, Doak L, Root J. Teaching Patients with Low Literacy Skills,. *American Journal of Nursing*. 1996;96(12):16M.

APPENDICES

Algorithms: Appendices A-F

Appendix A: Sample Management of Eclampsia Algorithm




Dr G Hilton and Colleagues (2012) - Lucile Packard Children's Hospital at Stanford

Committee Opinion No. 514. American College of Obstetricians and Gynecologists. Obstet Gynecol 2011;118:1465-8

Appendix B: Sample Treatment of Severe Preeclampsia Algorithm


Treatment for Severe Preeclampsia



Emergent therapy for acute-onset, severe hypertension with preeclampsia or eclampsia. Committee Opinion No. 514. American College of Obstetricians and Gynecologists. Obstet Gynecol 2011;118:1465-8

Appendix C: Suspected Preeclampsia Algorithm

Suspected Preeclampsia Flowchart
Diagnosis and Management




NOTE: There are no longer real differences in management between Preeclampsia and Gestational HTN in BP Management and decision to deliver. TREAT BP ACCORDINGLY and DELIVER for abnormal labs or symptoms

Appendix D: Preeclampsia Early Recognition Tool (PERT)


Preeclampsia Early Recognition Tool (PERT)

ASSESS	NORMAL (GREEN)	WORRISOME (YELLOW)	SEVERE (RED)
Awareness	Alert/Oriented	• Agitated/Confused • Drowsy • Difficulty speaking	• Unresponsive
Headache	None	• Mild headache • Nausea, Vomiting	• Unrelieved Headache
Vision	None	• Blurred or impaired	• Temporary blindness
Systolic BP (mmHG)	100-139	140-159	≥160
Diastolic BP (mmHG)	50-89	90-105	≥105
HR	61-110	111-129	≥130
Respiration	11-24	25-30	<10 or >30
SOB	Absent	Present	Present
O2 Sat (%)	≥95	91-94	≤90
Pain: Abdomen or Chest	None	• Nausea, Vomiting • Chest Pain • Abdominal Pain	• Nausea, Vomiting • Chest Pain • Abdominal Pain
Fetal Signs	• Category I • Reactive NST	• Category II • IUGR • Non-Reactive NST	• Category III
Urine Output (ml/hr)	≥50	30-49	≤30 (in 2 hrs)
Proteinuria	Trace	• +1/+2 • ≥300mg/24 hours	• >+3 • ≥5g/24hrs*
Platelets	>100	50-100	<50
AST/ALT	<70	>70	>70
Creatinine	<0.8	0.9-1.2	>1.2
Magnesium/Mag Sulfate Toxicity	• DTR +1 • Respiration 16-20	• Depression of patellar reflexes	• Respiration <12


*Level of Proteinuria is not an accurate predictor of pregnancy outcome



Appendix E: Eclampsia Algorithm



Appendix F: Evaluation and Treatment of Antepartum and Postpartum Preeclampsia and Eclampsia in the Emergency Department Part 1 AND 2



Evaluation and Treatment of Antepartum and Postpartum
 Preeclampsia and Eclampsia in the Emergency Department

TREATMENT	<p>1st Line Anti-Hypertensive Treatment: Labetalol & Hydralazine* Target BP: 140-160/90-100/90 (BP<140/90 = decreased fetal perfusion) See CMQCC Preeclampsia Toolkit for "Antihypertensives in Preeclampsia" for 2nd line therapy</p>	<p>Magnesium</p>	
	<p>LABELALOL as Primary Anti-Hypertensive</p> <ol style="list-style-type: none"> Administer Labetalol 20 mg IV Repeat BP in 10 min <ul style="list-style-type: none"> If BP threshold is still exceeded, administer labetalol If SBP<160 and DBP<100, continue to monitor closely Repeat BP in 10 min <ul style="list-style-type: none"> If BP threshold is still exceeded, administer labetalol 80 mg IV If SBP<160 and DBP<100, continue to monitor BP closely Repeat BP in 10 min <ul style="list-style-type: none"> If BP threshold is still exceeded, administer hydralazine 10 mg IV If SBP<160 and DBP<100, continue to monitor closely Repeat BP in 20 min; if BP threshold is still exceeded, obtain emergent consultation from maternal-fetal medicine, internal medicine, anesthesiology, or critical care Once target BP achieved, monitor BP q10 min for 1 hour, q 15 min for 2nd hour 	<p>HYDRALAZINE as Primary Anti-Hypertensive</p> <ol style="list-style-type: none"> Administer hydralazine 5 or 10 mg IV Repeat BP in 20 min <ul style="list-style-type: none"> If BP threshold is still exceeded, administer hydralazine 10 mg IV If SBP<160 and DBP<100, continue to monitor closely Repeat BP in 20 min <ul style="list-style-type: none"> If BP threshold is still exceeded, administer labetalol 20 mg IV If SBP<160 and DBP<100, continue to monitor BP closely Repeat BP in 10 min <ul style="list-style-type: none"> If BP threshold is still exceeded, administer labetalol 40 mg IV and obtain emergent consultation from maternal-fetal medicine, internal medicine, anesthesiology, or critical care If SBP<160 and DBP<100, continue to monitor closely Once target BP achieved, monitor BP q10 min for 1 hour, q 15 min for 2nd hour 	<p>Initial Treatment</p> <ol style="list-style-type: none"> Loading Dose: 4-6 gm over 15-20 min Maintenance 1-2 gm/hr Close observation for signs of toxicity <ul style="list-style-type: none"> Disappearance of deep tendon reflexes Decreased RR, shallow respirations, shortness of breath Heart block, chest pain Pulmonary edema
			<p>If Patient Seizes While on Magnesium:</p> <ol style="list-style-type: none"> Secure airway and maintain oxygenation Give 2nd loading dose of 2 gm Magnesium over 5 min If patient seizes after 2nd magnesium bolus, consider the following: <ul style="list-style-type: none"> Midazolam 1-2mg IV; may repeat in 5-10 min OR Lorazepam 2 mg IV-may repeat OR Diazepam 5-10 mg IV. May repeat q15 min to max of 30 mg Phenytoin 1 g IV over 20 min
		<p>Seizures Resolve</p> <ol style="list-style-type: none"> Maintain airway and oxygenation Monitor VS, cardiac rhythm/ECG for signs of medication toxicity Consider brain imaging for: <ul style="list-style-type: none"> Head trauma Focal Seizure Focal neurologic findings Other neurologic diagnosis is suspected 	

*Labetalol & Hydralazine recommendations based on 2011 ACOG Committee Opinion #514 and Practice Bulletin #33, Reaffirmed 2010

Simulations/Drills: Appendices G-N

Appendix G: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: General Information

Severe Preeclampsia and Eclampsia in LDR v2.0 SimMan3G

Part 1 – General Information

Authors: Mark Meyer MD, Darin Bowers MA – Southern California Permanente Medical Group

Scenario	SimMan3G – LDR Severe Preeclampsia & Eclampsia v2.0
Scenario Time	15-20 minutes
Debriefing Time	20-45 minutes – longer if used as 1 st scenario and requires more time to discuss non-technical skills (teamwork, communication, etc.) and CMQCC guidelines
Target Group	L&D nurses, OB physicians, Anesthesiologists, CRNA's, & scrub techs.
Case Summary	<p>This is a case of a patient on L&D who is being induced for mild preeclampsia. The patient develops severe preeclampsia and eclampsia that requires anti-hypertensive treatment as well as additional magnesium to control seizures. Despite maximal magnesium therapy, the patient continues to seize and the patient will require additional medications to control her seizures. In addition, the patient's SpO2 will fall due to airway occlusion during/after the seizure. Simple repositioning of the head and opening the airway will restore SpO2. No intubation is required, but this could be required if desired. This case is designed to ensure staff are following ACOG & CMQCC guidelines for appropriate treatment of preeclampsia and eclampsia. Therefore, there is a great emphasis on appropriate medication dosing and timing per these guidelines.</p> <p>It is critical that the participants recognize the patient is seizing. Unfortunately, the effectiveness of the SimMan3G seizure feature is limited, so confederate may be required to point out the seizure if the team does not recognize this. Fetal monitoring simulators can also be used, however, a non-reassuring fetal heart tracing may prompt the treatment team move the patient to the OR before the patient has received appropriate treatment for eclampsia and is stable for urgent c-section.</p>
Teaching Personnel	<ol style="list-style-type: none"> 1. GUI operator 2. Observer to note team communication and medical management skills – will serve as lead debriefer 3. Family member to voice observation of seizure signs if staff doesn't recognize seizure (essential if using SimMan Classic and SimMan3G) 4. OB physician for clinical expertise if the lead debriefer is not OB 5. Voice of patient – could be GUI operator
Participants	<ol style="list-style-type: none"> 1. 1-2 OB Physicians 2. 2-4 L&D nurses – varies depending on usual staffing on your unit 3. 1 CNM 4. Anesthesiologist and/or CRNA
Learning Objectives	<ol style="list-style-type: none"> 1. Demonstrate effective teamwork and communication skills with a focus on adequate shared mental model and role clarity. This includes clear identification of all team members and SBAR to new team members as they arrive. 2. Diagnose severe preeclampsia 3. Treat hypertension per CMQCC Preeclampsia/Eclampsia guidelines 4. Provide appropriate initial management of eclamptic seizures with magnesium 5. Manage eclamptic seizures when magnesium is ineffective 6. Maintain airway and oxygenation in seizing and post-ictal patient
References	<ol style="list-style-type: none"> 1. Emergent therapy for acute-onset, severe hypertension with pre- eclampsia or eclampsia. Committee Opinion No. 514. American College of Obstetricians and Gynecologists. Obstet Gynecol 2011;118: 1465–8 2. Preeclampsia care guidelines and compendium of best practices. California Maternal Quality Care Collaborative (CMQCC). 2012.

Used with permission, Kaiser Permanente and Mark Meyer, MD and Darin Bowers, MA

Appendix H: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: Learning Objectives

Severe Preeclampsia and Eclampsia in LDR v2.0 SimMan3G

Part 1 – General Information

Authors: Mark Meyer MD, Darin Bowers MA – Southern California Permanente Medical Group

Scenario	SimMan3G – LDR Severe Preeclampsia & Eclampsia v2.0
Scenario Time	15-20 minutes
Debriefing Time	20-45 minutes – longer if used as 1 st scenario and requires more time to discuss non-technical skills (teamwork, communication, etc.) and CMQCC guidelines
Target Group	L&D nurses, OB physicians, Anesthesiologists, CRNA's, & scrub techs.
Case Summary	<p>This is a case of a patient on L&D who is being induced for mild preeclampsia. The patient develops severe preeclampsia and eclampsia that requires anti-hypertensive treatment as well as additional magnesium to control seizures. Despite maximal magnesium therapy, the patient continues to seize and the patient will require additional medications to control her seizures. In addition, the patient's SpO2 will fall due to airway occlusion during/after the seizure. Simple repositioning of the head and opening the airway will restore SpO2. No intubation is required, but this could be required if desired. This case is designed to ensure staff are following ACOG & CMQCC guidelines for appropriate treatment of preeclampsia and eclampsia. Therefore, there is a great emphasis on appropriate medication dosing and timing per these guidelines.</p> <p>It is critical that the participants recognize the patient is seizing. Unfortunately, the effectiveness of the SimMan3G seizure feature is limited, so confederate may be required to point out the seizure if the team does not recognize this.</p> <p>Fetal monitoring simulators can also be used, however, a non-reassuring fetal heart tracing may prompt the treatment team move the patient to the OR before the patient has received appropriate treatment for eclampsia and is stable for urgent c-section.</p>
Teaching Personnel	<ol style="list-style-type: none"> 1. GUI operator 2. Observer to note team communication and medical management skills – will serve as lead debriefer 3. Family member to voice observation of seizure signs if staff doesn't recognize seizure (essential if using SimMan Classic and SimMan3G) 4. OB physician for clinical expertise if the lead debriefer is not OB 5. Voice of patient – could be GUI operator
Participants	<ol style="list-style-type: none"> 1. 1-2 OB Physicians 2. 2-4 L&D nurses – varies depending on usual staffing on your unit 3. 1 CNM 4. Anesthesiologist and/or CRNA
Learning Objectives	<ol style="list-style-type: none"> 1. Demonstrate effective teamwork and communication skills with a focus on adequate shared mental model and role clarity. This includes clear identification of all team members and SBAR to new team members as they arrive. 2. Diagnose severe preeclampsia 3. Treat hypertension per CMQCC Preeclampsia/Eclampsia guidelines 4. Provide appropriate initial management of eclamptic seizures with magnesium 5. Manage eclamptic seizures when magnesium is ineffective 6. Maintain airway and oxygenation in seizing and post-ictal patient
References	<ol style="list-style-type: none"> 1. Emergent therapy for acute-onset, severe hypertension with pre- eclampsia or eclampsia. Committee Opinion No. 514. American College of Obstetricians and Gynecologists. Obstet Gynecol 2011;118: 1465–8 2. Preeclampsia care guidelines and compendium of best practices. California Maternal Quality Care Collaborative (CMQCC). 2012.

Used with permission, Kaiser Permanente and Mark Meyer, MD and Darin Bowers, MA

Appendix I: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: Patient Background Information

Severe Preeclampsia and Eclampsia in LDR v2.0 SimMan3G

Part 3 – Patient Background Information

Authors: Mark Meyer MD, Darin Bowers MA – Southern California Permanente Medical Group

Patient Information and Background OPTION #1	
Age	28
Weight	182 lb
HPI	28 y/o G1P0 at 39 weeks admitted several hours ago after presenting to L&D triage with severe swelling BP was slightly elevated at 142/93. Patient was otherwise asymptomatic . Patient was given magnesium bolus and is currently on magnesium drip. Laboratory evaluation was WNL except for proteinuria..
PMHx	None
Medications	Prenatal Vitamins
Allergies	Sulfa
Social Hx	Married, sales person at Nordstrom's. No ETOH, Drugs, Tobacco.
Presentation	Patient calls the nurse after developing a moderate headache. The nurse should evaluate the patient's VS and note that she is now markedly hypertensive.
Vital Signs	Most recent set before nurse enters room: T 98.2 P 93 R 16 BP 142/93

Used with permission, Kaiser Permanente and Mark Meyer, MD and Darin Bowers, MA

**Appendix J: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G:
 Equipment/ Materials List**

Severe Preeclampsia and Eclampsia in LDR Unit v2.0 SimMan3G

Part 4 – Equipment/Materials List

Authors: Mark Meyer MD, Darin Bowers MA - Southern California Permanente Medical Group

Simulation Equipment:

Option #1 – SimMan 3G	
For video debriefing:	Webcam
	Additional laptop with Laerdal's Debrief Viewer software installed
	Digital Projector
	Flash drive to transfer Debrief Viewer files between laptops
Power strip	
Foam tape	

Patient Care Equipment:

ID band on SimMan3G
IV in place
IVF - 0.9NS or whatever IVF you prefer in this setting
BP cuff and pulse oximeter
100% O2 Non-Rebreather Mask
Adult ambu bag and oxygen tubing
Suction module, canister, tubing, yankauer tip
Stool for shoulder dystocia
Fetal monitor - OPTIONAL
Delivery room set up
Epidural setup – Optional
Adult crash cart that includes:
Airway equipment for intubation OPTIONAL
Defibrillator that can be used as cardiac monitor

Medications:

Magnesium 6g
Magnesium 4g
Magnesium 2g
Pitocin OPTIONAL
Labetalol
Hydralazine
Benzodiazepine – Ativan, Versed, or Valium

Used with permission, Kaiser Permanente and Mark Meyer, MD and Darin Bowers, MA

Appendix K: Severe Preeclampsia/Eclampsia IN LDR v2.0 SimMan 3G: Program Algorithm & GUI Notes

Severe Preeclampsia and Eclampsia in LDR Unit v2.0 SimMan3G

Part 5 - Program Algorithm & GUI Notes

<p>Patient: Monitor: OB</p> <p>Sinus Rhythm: 93 bpm Monitor Controls SpO2 = 98 % TPeri = 99.2 °F Respiration Rate: 12 CO2 Exhalation: Off Blood Pressure: 177/110 Handler Seizure airway: Start</p>	<p>Initial State</p> <ol style="list-style-type: none"> This is a case of a patient on L&D who is being induced for mild preeclampsia on a magnesium drip. The patient develops severe preeclampsia and eclampsia. Either 2 doses of hydralazine or 3 doses of labetalol will be required to control BP. Despite maximal magnesium therapy, the patient may require additional medications to control seizures. The case will end when the patient's BP and seizures are well controlled. There is a great emphasis on appropriate medication & timing to insure staff are following ACOG & CMQCC guidelines for preeclampsia & eclampsia. Since labetalol & hydralazine require 10-20 min for effect, the instructors will speed up the case by announcing 10 minutes has passed for each minute after labetaol or hydralazine is given. See frames below for details. The SimMan 3G GUI operator will be the voice of the patient & will use the seizure feature on this simulator. Since the seizure feature has limited realism, it is recommended that a confederate notify the team that the patient is having a seizure if the treatment team does not recognize this. In this frame, the patient calls the nurse because she has a moderate headache. The responding nurse should note the VS & report the BP to the physician or CNM that responds to the call for help Once the physician or CNM orders either labetaol or hydralazine, or vocalizes the diagnosis of severe preeclampsia, then click "Advance next frame." In the next frame, the patient begins to seize for the first time. NOTE: The "Seizure airway" handler – this will restore SpO2 to 98% when the airway is opened after the patient seizes
<p>Call for physician Call for additional nurse Advance to next frame ORDER Labetalol ORDER Hydralazine Dx of Severe Preeclampsia</p>	
<p>Eyes Eyes half open Blinking speed: Off Convulsions: Tonic-Clonic Fluids & Secretions Froth: On Airway Trismus: On Start Trend: Seizure desaturation</p>	<p>SEIZURE #1</p> <ol style="list-style-type: none"> The patient is now seizing. When using SimMan3G, the seizures are most notable if the arms are out from the body & visible to the participants. If using the bodily fluid function of the simulator, the patient will foam at the mouth as well. Note the trend: "Seizure desaturation" which causes the SpO2 to fall into the 80's. The team should reposition the patient's head & keep the airway open, which will restore the SpO2 to normal levels. The team should be working to administer an antihypertensive & an additional magnesium bolus. The seizure will last 60 sec & the scenario will advance automatically. Click "Advance next frame" to advance more quickly if desired.
<p>Check airway patency Open airway Advance to next frame FT=1:00 ORDER Magnesium bolus Oxygen</p>	
<p>Convulsions: None Eyes Blinking speed: Normal Fluids & Secretions Froth: Off Airway Trismus: Off Blood Pressure: 179/109</p>	<p>SEIZURE #1 STOPS</p> <ol style="list-style-type: none"> The seizure stops & the patient is post-ictal. She will respond to simple questions & commands, but will have somewhat slurred speech. The patient must appear awake enough to protect her airway or the team may try to intubate the patient. If the team has not already done so, they must reposition the head and open the airway to restore a normal SpO2! The scenario advances when either labetalol or hydralazine is given. Alternatively click "Advance next frame" to advance. The team should be administering an additional magnesium bolus at this time.
<p>2g Magnesium GIVEN 10mg Hydralazine GIVEN Advance to next frame 20mg Labetalol GIVEN 5mg Hydralazine GIVEN 4g Magnesium GIVEN</p>	
<p>Blood Pressure: 175/100</p>	<p>10 Min Since Anti-HTN and Mag Bolus</p> <ol style="list-style-type: none"> At approximately one minute into this frame, the instructor should call out that 10 minutes have passed which should prompt an additional BP check. If labetalol was given initially, another dose is indicated. The scenario will advance when the next dose of labetalol is given. If hydralazine was initially given, click "Advance next frame" as it is given q20 min prn.
<p>40mg Labetalol GIVEN Advance to next frame Click to add learner event</p>	
<p>Convulsions: None Fluids & Secretions Froth: Off Eyes Blinking speed: Normal Airway Trismus: Off Blood Pressure: 167/100</p>	<p>20 min</p> <ol style="list-style-type: none"> In this frame, approximately 20 min have passed since HTN meds and magnesium were given. At approximately one minute into this frame, call out that an additional 10 minutes have passed. This should prompt an additional BP check. The BP has improved but requires another dose of labetalol or hydralazine. The instructor should also call out that the mag bolus is complete. Click on either medication or "Advance next frame" and the case will end. If desired, click "Advance to Seizure #2" and the patient will seize again.
<p>Advance to Seizure #2 10mg Hydralazine GIVEN Advance to next frame 80mg Labetalol GIVEN</p>	
<p>Eyes Eyes half open Blinking speed: Off Convulsions: Tonic-Clonic Fluids & Secretions Froth: On Airway Trismus: On Start Trend: Seizure desaturation (Start: 0 min)</p>	<p>SEIZURE #2</p> <ol style="list-style-type: none"> The patient is seizing despite maximal magnesium therapy & should prompt the team to use other meds to control the seizure. The team should consider benzodiazepines or dilantin. The seizure lasts minute & go back to the previous frame to insure that the additional dose of labetalol or hydralazine is given.
<p>Check airway patency Check breathing Advance to next frame FT=1:00 ORDER Benzo Benzo GIVEN</p>	<p>END</p> <ol style="list-style-type: none"> At approximately one minute into this frame, call out that 10 minutes have passed. The BP & seizures are now controlled. If indicated, the patient is now stable to go to the OR for c-section.
<p>20 min 20 min</p>	<p>Click to add learner event Click to add learner event Click to add learner event</p>

Used with permission, Kaiser Permanente and Mark Meyer, MD and Darin Bowers, MA

Appendix L: Severe Preeclampsia/Eclampsia In LDR v2.0 SimMan 3G: Debriefing Objectives

Severe Preeclampsia and Eclampsia in LDR v2.0 SimMan3G

Part 6 – Debriefing Objectives

Authors: Mark Meyer MD, Darin Bowers MA – Southern California Permanente Medical Group

Scenario	v2.0 LDR Severe Preeclampsia_Eclampsia.sce
Est. Debriefing Time	20-30 minutes
Debriefing Objectives (4-5 Max)	NON-TECHNICAL SKILLS
Non-Technical	<p>SHARED MENTAL MODEL - "Does everyone know what's going on?"</p> <ol style="list-style-type: none"> SBAR responding team members to establish Shared Mental Model <ol style="list-style-type: none"> Stituation/Background – suggested components required – Age, Gravida, Parity, Gestational age, VS with emphasis on BP, allergies, what interventions/treatments have already been given, pertinent symptoms, and any contraindications to treatment – in this case, asthma would be contraindication for using labetalol Assessment – What is our assessment of this patient's condition? e.g. "This patient has severe preeclampsia" Recommendations <ol style="list-style-type: none"> Are these consistent with CMQCC recommendations? Initial recommendations don't have to be comprehensive, but rather, adequate to generate and share a general treatment plan. Details (e.g.dosing) can come later Sample: "We need to treat the patient's hypertension, monitor VS and EFM closely, draw labs call for anesthesia, and be prepared if the patient begins to seize." Commonly, team may only receive Situation and Background from the nurse. Focus on encouraging the nursing staff to give their assessment and recommendation Did entire team and/or latecomers get adequate report? Team Leader(s) give SBAR back to team after initial assessment is complete <ol style="list-style-type: none"> Critical as first responder often does not provide assessment and plan. This can be due to limited information, but often this is medical cultural issue i.e. not all team members are comfortable speaking up. Provides clear plan to team and provides opportunity for team feedback
Medical Management	<p>ROLE CLARITY - only complete if it includes "Task Clarity as well"</p> <ol style="list-style-type: none"> Clear Role Assignment <ol style="list-style-type: none"> Leader(s) – often a physician/nurse team form the most effective leadership Did personnel identify themselves? e.g. OB physician, midwife, nurse, anesthesia, etc. Primary nurse assigns tasks vs. "self selection" of roles/tasks "Task Clarity" - clear verbalization of who is doing what task(s). <ol style="list-style-type: none"> The tasks that must be done are often not clearly assigned as part of the team member's role. e.g. There are several nursing tasks, but it may not be clear who is doing each. Regardless of how roles and tasks are assigned, these must be verbalized clearly so that entire team knows who is responsible for necessary task <p style="text-align: center;">MEDICAL MANAGEMENT SKILLS</p> <ol style="list-style-type: none"> Treat hypertension per CMQCC guidelines <ol style="list-style-type: none"> Initial treatment given <30 minutes is CMQCC goal Appropriate drugs, dosing, indications and timing for repeated dosing? Eclampsia/ Seizure <ol style="list-style-type: none"> Magnesium given in timely fashion? Additional mag given if necessary? Medications for treatment of seizures refractory to magnesium – E.g. benzodiazepines. Maintain airway & oxygenation in seizing and post-ictal patient See CMQCC guidelines for more details
1. Shared Mental Model	
2. Role Clarity	
3. Hypertension	
4. Seizures	
5. Airway/oxygenation	

permission, Kaiser Permanente and Mark Meyer, MD and Darin Bowers, MA

Used with

Appendix M: Severe Preeclampsia/Eclampsia In LDR v2.0 SimMan 3G: Debriefing Guide/Evaluation

Part 7 Debriefing Guide/Evaluation – Make notes & time of notable behavior for use during video debriefing

<p>Shared Mental Model/SBAR</p> <ol style="list-style-type: none"> Nurse to team responding to call for help <ol style="list-style-type: none"> Are all elements present? What elements are missing, why? Did the responding team “receive” the report? <ol style="list-style-type: none"> Report given only to physician? Team doing other tasks and not listening? Team leader gather team for report? Report given to team members who arrive later? Does physician give SBAR back to team after initial assessment? 	<p>Shared Mental Model/SBAR Notes</p> <ol style="list-style-type: none"> SBAR elements? Did team “receive” report? New team members get report? Physician gives SBAR back to team?
<p>Role Clarity</p> <ol style="list-style-type: none"> Clear nursing and physician leader? Other Roles clearly established? <ol style="list-style-type: none"> Self-assignment Team leader assigns roles/tasks Are roles clearly verbalized to ALL team members? 	<p>Role Clarity Notes</p> <ol style="list-style-type: none"> Clear nursing & physician leader? Roles clearly established? Roles clearly verbalized?
<p>Situational Awareness & Assertion</p> <ol style="list-style-type: none"> Ongoing monitoring & crosschecking? (Assess condition & response to treatment) Recognizes critical information in timely fashion i.e. “The Gorilla” Effective assertion to appropriate team member? e.g. 2 challenge rule, etc. 	<p>Situational Awareness & Assertion Notes</p> <ol style="list-style-type: none"> Monitoring of situation? Recognizes critical information Effective assertion?
<p>Closed Loop Communication</p> <ol style="list-style-type: none"> Responds appropriately to input from team? (leaders open to input, rationale for actions?) Readback of orders 	<p>Closed Loop Communication Notes</p> <ol style="list-style-type: none"> Responds to input? Readback of orders
<p>Cognitive Skills/ Medical Management</p> <ol style="list-style-type: none"> Critical medical management actions (varies with each scenario) Considers other diagnoses when patient doesn’t respond as expected Avoids medication errors 	<p>Cognitive Skills/ Medical Management Notes</p> <ol style="list-style-type: none"> Critical medical management actions? Considers other diagnoses when appropriate? Avoids medication errors?
<p>Psychomotor Skills/Equipment Competency</p> <ol style="list-style-type: none"> Quickly locates critical equipment Competent with critical equipment/systems 	<p>Psychomotor Skills/Equipment Competency Notes</p> <ol style="list-style-type: none"> Locates critical equipment Competent with equipment

Used with permission, Kaiser Permanente and Mark Meyer, MD and Darin Bowers, MA

Appendix N: Simulation Scenario: Hypertension in Pregnancy, Help with Seizure

Simulation: HELLP with Seizure

Leslie Cragin, CNM
Ana Delgado, CNM
Ocean Berg, RN, MSN, IBCLC, UC San Francisco, OB/GYN Department

Topic: Hypertension in Pregnancy Scenario: HELLP with seizure

Duration of Scenario: 6 - 13 min

General Description of the Scenario:

Jackie is a 17 yo G1P0 @ 36 weeks by sure normal LMP. She came to triage accompanied by her sister after beginning uterine contractions 8 hours ago, which have been increasing in intensity and frequency. The contractions are every 3-4 minutes, lasting a minute. She is here with her older sister. Jackie complains of a strong headache beginning 2 days ago. The baby is moving less than before labor began. She was admitted for labor 4 hours ago with V/E 4cm, 70%, -2 station.

Brief Medical/OB History:

- Regular visits, no chart available
- Fundal height = 34
- Admit labs: hgb 9.2, hct 30, platelets 90,000

Objectives:

Cognitive:

1. Accurately identify risk factors for severe preeclampsia/HELLP
2. Identify the differential diagnosis for eclampsia
3. Identify medications to be used in managing an eclamptic seizure
4. Know the steps in management eclampsia

Technical:

1. Provide protection from injury and patent airway during seizure
2. Evaluate for interval to delivery
3. Evaluate fetal status
4. Prepare for fetal resuscitation and potentially postpartum hemorrhage

Behavioral:

1. Calls for help in a timely manner
2. Communication with team
3. Maintains a calm demeanor during the emergency
4. Clear communication with the frightened family members

Roles of the participants:

RN, CNM, extra RN, obstetrician,

- Facilitator taps out Fetal heart rate
- MD is slow to come in after being called– doesn't intervene or direct but does ask what is happening

Roles of the Confederates:

Patient in PartoPants®, Significant other

Equipment: Partopants, bed, sheets, footstool, baby, IV pole/set/fluid, Doppler or fetoscope, delivery set,

Simulator: Actress with PartoPants

Opening scene: Mother is laboring with _____

Progression of Scenario

Time	Events for Actress and Confederates	Appropriate Actions	Symptoms/Results of inappropriate action
0-5 min	<p>Patient (IV in place); midwife/FCM; RN and significant other in room;</p> <p>Patient in labor with ctx Q 3 min</p> <p>FHR 120's</p> <p>Pt begins to seize at about 5 minutes into scenario - seizure lasts 90 seconds</p> <p>Fetal bradycardia to 80 BPM for 3 min begins with seizure and lasts 3 min</p>	<ul style="list-style-type: none"> • Cleans hands <p>Begins assessment: talks with patient</p> <p>Requests vital signs</p> <p>Asks about urine and proteinuria</p> <p>May ask for additional labs</p> <p>May turn mom into side-lying</p> <p>Notes FHR</p>	<p>Initial vitals 138/89</p> <p>P=110</p> <p>No pain meds given yet</p> <p>No proteinuria</p> <p>V/E if done 8cm, 100 % 0 station</p> <p>BP stays in this range - never severely elevated</p>
4/5-10	<p>Seizure resolves</p> <p>Sister asks what is happening</p> <p>FHR 160's then back to normal</p>	<p>Pt turned to L side, O2 on</p> <p>Mag. sulfate ordered: 4-6 gm IV over 15 minutes or 5 gm IM in each buttock if no IV</p> <p>NO BP meds since BP is not elevated</p> <p>Calls for help</p> <p>Evaluates FHR</p> <p>Gives accurate concise report to attending</p>	<p>If no Mag. ordered by 2 min postictal, another seizure begins- this should be treated with MgSO4, diazepam ok, but NOT optimal</p>

	Pt is postictal/sleepy Pt involuntarily pushes	V/E	STOP SCENARIO
--	---	-----	---------------

Guide for review of simulation:

(Remember to focus on cues from the video, these are only triggers for discussion.)

General:

1. How did that feel?
2. Would someone give an overview of the scenario?
3. What did you see?
4. What went well?
5. What didn't?
6. Was there anything in the **10 commandments** that would've helped you?

Cognitive:

1. What were you thinking when you heard about the report?
2. What are the risk factors for pre-eclampsia/severe and HELLP?
3. What are the signs and symp of HELLP?
4. What labs would help to evaluate this pt?
5. What other emergencies/complications follow eclampsia (PPH, neonatal compromise)?

Technical:

1. What should be done to protect the patient?
2. What are the components of intrauterine resuscitation?
3. What are the medications to be used in eclampsia with severe HTN?

Behavioral: Focus on 2-3 points

1. Know your environment and team
2. Plan and anticipate
3. Assume the role of leader
 - a. Who was the leader?
 - b. How did that go? (ask leader and participants)
4. Communicate in an effective manner with the team, the patient and her family
 - a. How was the interaction between the midwife/FCM and nurse?
 - b. How was the communication with the patient?
5. Delegate appropriately
6. Allocate attention wisely
7. Use all your available resources
8. Use all your available information
9. Call for help in a timely manner
 - a. What made you call for help?
10. Maintain professional conduct/attitude at all times.

Used with permission of Leslie Cragin, CNM, Ana Delgado, CNM, Ocean Berg, RN,
MSN, IBCLC, UC San Francisco, OB/GYN Department

Patient Education: Appendices O-Q

Appendix O: Preeclampsia Foundation Signs and Symptoms Information Sheet

Ask Your Doctor or Midwife

Preeclampsia

What Is It?

Preeclampsia is a serious disease related to high blood pressure. It can happen to any pregnant woman.

Risks to You

- Seizures
- Stroke
- Organ damage
- Death

Risks to Your Baby

- Premature birth
- Death

Signs of Preeclampsia




Stomach pain




Headaches



Feeling nauseous; throwing up



Seeing spots



Swelling in your hands and face



Gaining more than 5 pounds in a week

What Should You Do?

Call your doctor right away. Finding preeclampsia early is important for you and your baby.

For more information go to www.preeclampsia.org

Copyright © 2010 Preeclampsia Foundation. All Rights Reserved.

To order materials, go to: www.preeclampsia.org/market-place

Appendix P: SAMPLE: Discharge Instructions Following Delivery with Diagnosed Preeclampsia

[NAME OF HOSPITAL, CLINIC OR PRACTICE]
[PHONE NUMBER]

DISCHARGE INSTRUCTIONS FOLLOWING DELIVERY OF BABY PREECLAMPSIA

During your hospitalization, you have been treated for preeclampsia or HELLP syndrome. Preeclampsia is a problem that can occur in the late stages of pregnancy and even *during the first few weeks postpartum* (after delivery of your baby), and causes high blood pressure, protein in the urine and sometimes other symptoms such as headaches, blurred vision, breathlessness, and swelling of the hands or face. In the past, it has been called “toxemia” or “pregnancy-induced hypertension”. HELLP syndrome is a variation of preeclampsia that directly affects your liver and blood platelets.

Preeclampsia can be mild or severe. If it isn’t treated, preeclampsia can turn into a serious problem called “eclampsia” in which seizures occur.

When you go home, follow these instructions:

- Keep follow-up appointments with your doctor. These may be very frequent and are very important for your health.
- Take all medications prescribed for you exactly as ordered.
- Weigh yourself at the same time each day. Write down your weight and take this record with you to your doctor visits.
- If ordered by your doctor, monitor your blood pressure at home.
- Ask your doctor if you need to check your urine at home for protein.
- Eat a healthy, balanced diet. Your doctor will tell you if you need to follow any special restrictions in what you eat.
- Don’t smoke.
- Don’t drink alcohol or use any drugs not prescribed to you.
- Ask your doctor before taking any medications that he or she didn’t prescribe for you. This includes any over-the-counter medications.

Call your doctor if:

- Your blood pressure is greater than _____ systolic (the top or first number).
- Your blood pressure is greater than _____ diastolic (the bottom or second number).
- You have a severe headache or dizziness.
- You have any headache that is not relieved with Tylenol or ibuprofen (e.g., Advil™, Motrin™).
- You have pain in your belly, especially the right upper area below your ribs.
- You have blurry or double vision, see spots or auras.
- Your swelling is worse.
- You gain more than 3 pounds in 3 days.
- You have serious difficulty catching your breath.
- You have any new or unusual symptoms.
- You have any questions or concerns.

If you have any of the above symptoms, call [phone number] immediately. If you are unable to reach your physician you need to go to the emergency room for evaluation. Be sure to tell them you just had a baby and you had preeclampsia.

Additional information about preeclampsia can be found at www.preeclampsia.org. There you will find accurate information about preeclampsia and related disorders of pregnancy, as well as a very friendly and helpful community of women with whom you can discuss any concerns or questions.

Localization (where indicated), photocopying and usage is recommended by the Preeclampsia Foundation © 2011

Used with permission from the Preeclampsia Foundation

Appendix Q: SAMPLE: Discharge Sheet for Preeclampsia, Eclampsia and HELLP Syndrome Patients

**Discharge Information for Patients with Diagnosis of Preeclampsia,
HELLP Syndrome or Eclampsia**

Your Medications include the following:

- 1) _____ To be taken every ____ hours.
- 2) _____ To be taken every ____ hours.
- 3) _____ To be taken every ____ hours.

Your postpartum follow-up appointment has been made with Dr. _____ in ____ days.

You have been instructed to check your blood pressure at home daily: Yes____ No____
Call your physician if your blood pressure is greater than _____ systolic (top number)
and/or

If your blood pressure is greater than _____ diastolic (bottom number)

Call your physician if:

- Your temperature is greater than 100.4.
- Your bleeding is greater than a heavy menses.
- You have any headache that is not relieved with Tylenol or ibuprofen (e.g., Advil™, Motrin™).
- You have pain in your belly, especially the upper area below your ribs.
- You have blurry or double vision, see spots or flashing lights.
- Your swelling is worse.
- You gain more than 3 pounds in 3 days.
- You have serious difficulty catching your breath.
- You have any new or unusual symptoms.
- You have any questions or concerns.

Miscellaneous: Appendices R-W

Appendix R: Prenatal Care Monitoring: Documentation of Diagnosis and Accurate ICD-9 Coding

Sarah Kilpatrick MD, Cedars Sinai Medical Center

Diagnostic accuracy

A large proportion of preeclampsia related deaths are preventable.^{1,2} Of preventable deaths and near-miss morbidities a large proportion of the preventable factors are provider related including delay in diagnosis, and incomplete or inappropriate management.^{3,4} Making a timely and correct diagnosis is fundamental for improving outcomes in preeclampsia. Further, an inaccurate diagnosis of preeclampsia is not uncommon. Only 45% of the ICD-9 diagnoses of mild preeclampsia were confirmed by chart review.⁵ Half of these were gestational hypertension and the other half severe preeclampsia. This study illustrates that even in a university hospital there is significant confusion around the classification of hypertension in pregnancy. If the diagnosis is not clear or is inaccurate (i.e. she is diagnosed with mild preeclampsia but she actually has severe preeclampsia), it is not surprising that she may receive the wrong treatment (e.g., expectant management rather than delivery). The need for clear diagnostic accuracy is even more important in a progressive disease, such as preeclampsia, that will typically worsen as the pregnancy progresses and correspondingly requires changes in management. Because the treatment will change as the disease evolves and or the gestational age advances, the provider must appropriately reevaluate the patient to capture the evolution of the disease. The correct diagnosis based on signs and symptoms for gestational hypertension, mild preeclampsia, severe preeclampsia, eclampsia, or superimposed preeclampsia must be made and appropriately documented. All interventions should be based on diagnosis and gestational age of the pregnancy.

Recommendations:

1. Clearly document and communicate initial diagnosis based on initial signs and symptoms.
2. Clearly document and communicate initial treatment plan, and plan for reevaluation based on initial diagnosis and gestational age.
3. Clearly document and communicate any changes in diagnosis and plan.
4. Hospital coders should review progress notes to identify most advanced stage of preeclampsia during hospital admission to code for discharge diagnosis.

Table 1: ICD-9 Codes for Hypertensive Diseases in Pregnancy and Postpartum

ICD-9 Code	Description
642.0	Benign essential hypertension complicating pregnancy, childbirth, and the puerperium
642.1	Hypertension secondary to renal disease, complicating pregnancy, childbirth, and the puerperium
642.2	Other pre-existing hypertension complicating pregnancy, childbirth, and puerperium
642.3	Transient hypertension of pregnancy (gestational hypertension)
642.4	Mild or unspecified preeclampsia
642.5	Severe preeclampsia
642.6	Eclampsia
642.7	Preeclampsia or eclampsia superimposed on preexisting hypertension

5th digits are used to denote the current episode of care (e.g. delivered, antepartum, postpartum)

Table 2: ICD-10 Codes for Hypertensive Diseases in Pregnancy and Postpartum

ICD-10 Code	Description
010.01	Preexisting essential hypertension complicating pregnancy
010.02	Preexisting essential hypertension complicating childbirth
010.03	Preexisting essential hypertension complicating the pureperium
011	Preexisting hypertension with preeclampsia
013	Gestational hypertension without significant proteinuria
014	<i>Preeclampsia [excludes preexisting hypertension with preeclampsia (011)]: see specific codes below:</i>
014.0	Mild to moderate preeclampsia
014.1	Severe preeclampsia [excludes HELLP syndrome (014.2)]
014.2	HELLP syndrome (HELLP)
014.9	Unspecified preeclampsia
015.0	Eclampsia in pregnancy
015.1	Eclampsia in labor
015.2	Eclampsia in the puerperium
015.9	Eclampsia, unspecified as to time period

Additional digits are used to denote the trimester (1, 2, 3, and 0=unspecified)

Suggested Strategies for Data Quality Improvement Opportunities

What strategies can be used to improve the quality of coding of hypertensive diseases in pregnancy? The release of new definitions for preeclampsia will create an educational opportunity that should be embraced. Small posters with the new definitions (and codes) should be placed on Labor and Delivery units and shared with the hospital coders. As a follow-up to a Grand Rounds on Preeclampsia, diagnostic criteria can be reinforced by going over a few cases at serial department meetings and asking the physicians to select the “right” diagnostic category. Finally, as a QI effort, as cases are reviewed in Perinatal Quality review Committees, attention can be focused on the correct categorization. Feedback can be quite effective.

REFERENCES

1. Isler CM, Rinehart BK, Terrone DA, Martin RW, Magann EF, Martin JN, Jr. Maternal mortality associated with HELLP (hemolysis, elevated liver enzymes, and low platelets) syndrome. *American Journal of Obstetrics and Gynecology*. Oct 1999;181(4):924-928.
2. Berg CJ, Harper MA, Atkinson SM, et al. Preventability of pregnancy-related deaths - Results of a state-wide review. *Obstetrics and Gynecology*. Dec 2005;106(6):1228-1234.
3. Geller SE, Rosenberg D, Cox SM, et al. The continuum of maternal morbidity and mortality: Factors associated with severity. *American Journal of Obstetrics and Gynecology*. Sep 2004;191(3):939-944.
4. Lawton BA, Wilson LF, Dinsdale RA, et al. Audit of severe acute maternal morbidity describing reasons for transfer and potential preventability of admissions to ICU. *Aust N Z J Obstet Gynaecol*. Aug 2010;50(4):346-351.
5. Geller SE, Ahmed S, Brown ML, Cox SM, Rosenberg D, Kilpatrick SJ. International Classification of Diseases-9th revision coding for preeclampsia: how accurate is it? *American Journal of Obstetrics and Gynecology*. Jun 2004;190(6):1629-1633; discussion 1633-1624.

Appendix S: Sample Preeclampsia/Eclampsia Medication Toolbox List

(Each institution should prepare its own medication toolbox)

Contents of Labor and Delivery Severe Preeclampsia and Eclampsia Box	
Drug contents	
Medications	Quantity
Magnesium sulfate 20g/500ml bag	1
Calcium gluconate 1000mg/10ml vial	1
Labetalol 100mg/20ml vial	1
Hydralazine 20mg/ml vial	1
Esmolol 100mg/10ml vial	1
Sodium Chloride 0.9% (NS) 10 ml vial	1
Propofol 10mg/ml, 20ml vial	
Supply contents	
Supplies	Quantity
20 ml syringe	1
10 ml syringe	2
3 ml syringe	2
18G needle (pink)	5
Alaris Pump Tubing Set	1

Kindly used with permission of Stanford University Medical Center and Gillian Hilton, MD 2013

Appendix T: Special Considerations: Amphetamine/Cocaine Drug abuse and hypotension

Mark Zakowski, MD
 Cedars Sinai Medical Center

Drug abuse is very common in women of childbearing age, with 12-21% of women 18-35 years using illicit drugs¹, especially cocaine and methamphetamine.² Many drugs not only affect maternal and fetal well-being, but also alter maternal responses to medications and treatment. In particular, acute and chronic amphetamine or cocaine use may result in physiologic and pharmacodynamic changes that prove difficult to manage in the parturient.

Cocaine blocks the presynaptic reuptake of sympathomimetic neurotransmitters including dopamine, serotonin and norepinephrine. Thus, the acute use of cocaine results in the relative neurotransmitter concentrations increasing at the catecholamine's site of action, producing adrenergic stimulation both centrally and peripherally. Cocaine may cause intense vasoconstriction producing hypertension, coronary ischemia, and reduced uterine artery blood flow; other effects include tachycardia, arrhythmias, altered sensorium, and hyperthermia. With chronic abuse of cocaine the sympathomimetic neurotransmitters become depleted and hypotension and lethargy may ensue.

Common treatment of cocaine/amphetamine induced hypertension and tachycardia may include the use of hydralazine and labetalol.³ However beta-blockers, including labetalol (beta-adrenergic to alpha-adrenergic blocking 7:1 when administered IV⁴) may produce unopposed alpha-adrenergic stimulation with worsening of coronary and peripheral vasoconstriction.⁵ Indeed, phentolamine may be the drug of choice for cocaine/amphetamine-induced hypertension.

Treatment of hypertension in the patient with chronic cocaine/amphetamine abuse may cause an exaggerated decrease in blood pressure. Hypotension may be difficult to treat due to altered vasopressor response and depleted endogenous catecholamine stores. Unexpected, severe hypotension may also occur after regional anesthesia or general anesthesia. Also noteworthy, ketamine administration may exacerbate symptoms of acute cocaine use, due to ketamine's sympathetic nervous system stimulation.

Amphetamines indirectly stimulate the sympathetic nervous system, causing the release of catecholamines from presynaptic vesicles. The symptoms are similar to cocaine intoxication and may include hypertension, arrhythmias, tachycardia, dilated pupils, hyperreflexia and even hyperthermia. Use of regional anesthesia may be associated with unpredictable hypotension, resistant to treatment. Cardiac arrest has also been reported following both regional and general anesthesia.

Parturients chronically using either cocaine or amphetamines are in a state of catecholamine depletion and potentially altered adrenergic receptor responses. Trauma patients may also exhibit catecholamine depletion, having maintained their blood pressure through extensive release of stored catecholamines. The initial antihypertensive treatment of elevated blood pressure due to acute intoxication with cocaine/amphetamine or even pain with coincident significant trauma may exacerbate subsequent hypotension.

Treatment of hypotension may be difficult in the catecholamine-depleted patient.

Ephedrine’s mechanism of action includes a direct effect as well as a significant secondary release of norepinephrine, which would be decreased in the catecholamine-depleted state, considerably blunting ephedrine’s effectiveness. The vasopressors of choice should be direct acting agents, administered intravenously. Preferred agents include phenylephrine, epinephrine and norepinephrine. While norepinephrine has the strongest direct acting vasoconstriction, it should be administered via central access. Please note that greater than typical bolus doses may be required. Infusions may also be very useful, depending on circumstances. Typical doses of direct acting vasopressors are listed below:

Phenylephrine 100 mcg IV bolus

Epinephrine 50-100 mcg IV bolus (stronger)

Norepinephrine 4-8 mcg IV bolus or infusion 2-20 mcg/min⁶ (risk of skin ischemia if given through small peripheral IV, prefer central line or secondarily large bore IV fast flowing)

Note that the same phenomenon of hypotension ‘resistant’ to vasopressor treatment has been reported following double calcium channel blocker usage – i.e., magnesium sulfate plus nifedipine. The physiologic difficulty with dual calcium channel blockers is post-depolarization low intracellular calcium. Normally the rise in intracellular calcium triggers a much greater release of calcium from the sarcoplasmic reticulum, causing contraction of the muscle cell. In the setting of post-dual calcium channel blocker hypotension, vasopressors may not be effective, because of calcium depletion. Fortunately, treatment with calcium (1 gram CaCl₂ IV) may reverse the primary problem – low intracellular calcium.

REFERENCES

1. National Center for Health Statistics. Health, United States, 2011: With Special Feature on Socioeconomic Status and Health. Hyattsville, MD. 2012
2. O’Brien CP, (2011). Chapter 24. Drug Addiction. In L.L. Brunton, B.A. Chabner, B.C. Knollman (Eds), *Goodman & Gilman’s The Pharmacological Basis of Therapeutics*, 12e.
3. Kuczkowski KM. Anesthetic Implications of drug abuse in pregnancy. *Journal of Clinical Anesthesia* 2003;15:382-394.

4. Morgan, JR. GE, Mikhail MS, Murray MJ. (2006). Chapter 12. Adrenergic Agonists & Antagonists. In G.E. Morgan, Jr., M.S. Mikhail, M.J. Murray (Eds), *Clinical Anesthesiology*, 4e.
5. Prosser JM, Perrone J. (2011). Chapter 181. Cocain, methamphetamine, and other amphetamines. In J.E. Tintinalli, J.S. Stapczynski, D.M. Cline, O.J. Ma, R.K. Cydulka, G.D. Meckler (Eds), *Tintinalli's Emergency Medicine: A Comprehensive Study Guide*, 7e.
6. Morgan, Jr. GE, Mikhail MS, Murray MJ. Chapter 12. Adrenergic Agonists & Antagonists. In: Morgan, JR. GE, Mikhail MS, Murray MJ, eds. *Clinical Anesthesiology*. 4th ed. New York: McGraw-Hill; 2006.

Appendix U: Sample Nursing Management Policy and Procedure

Nursing Management of Preeclampsia Sample Policy and Procedure

Brenda Chagolla, RN, MSN, CNS, University of California Davis Medical Center
Ocean Berg, RN, MSN, CNS, Nurse Family Partnership Program, San Francisco
Kristi Gabel, RNC-OB, C-EFM, MSN, CNS, Sutter Roseville Medical Center

PURPOSE:

To outline the nursing management of in patients who have preeclampsia including special considerations for management of patients on magnesium sulfate, patients on antihypertensive medications and management of eclampsia.

BACKGROUND:

Preeclampsia is a hypertensive disorder of pregnancy characterized by vasospasm and endothelial damage which may impact the cardiovascular, renal, hematological, neurologic, and hepatic systems as well as the uteroplacental unit. It is of unknown etiology. Preeclampsia is characterized by new onset of hypertension and proteinuria after 20 weeks gestation in a previously normotensive woman.

- Hypertension: two blood pressure reading of >140 systolic OR >90 diastolic taken at least six hours apart
- Proteinuria: 0.3gm of protein in a 24 hour urine collection

REPORTABLE CONDITIONS:

Notify provider for:

1. Repeated blood pressure greater than 160 systolic OR greater than 105 diastolic (taken at least 15 minutes apart).
2. New or worsening complaint of any of the following:
 - a. Headache
 - b. Visual changes
 - c. Right Upper Quadrant (RUQ) or epigastric pain
3. Abnormal lab values

ADMISSION:

1. Assess for absence or presence of:
 - a. Headache
 - b. Visual changes
 - c. Right upper quadrant or epigastric pain
 - d. Nausea/vomiting
 - e. General malaise

2. Assess upper or lower deep tendon reflexes.
3. Auscultate for lung sounds, noting any presence of rales, rhonchi, wheezing, etc.
4. Assess for generalized edema and significant, rapid weight gain.
5. Assess blood pressure using an appropriately sized blood pressure cuff with patient sitting or in the upright position with the patient's arm at the level of the heart. Do not reposition the patient to her left side and retake blood pressure. It will give a false lower reading.
6. Apply external fetal monitor (if viable fetus).
7. Prepare to obtain IV access as ordered by provider.
8. Prepare to administer medications to lower blood pressure and prevent seizure activity.
9. Prepare to monitor intake and output.
10. Maintain activity as ordered by provider. If on bedrest, maintain side-lying position as much as possible, avoiding supine position, and change position every two hours or more often as needed.
11. Provide emotional support and opportunity for patient family to verbalize questions, concerns and/or fears.
12. Assess maternal vital signs including: blood pressure as described above, respiratory rate, heart rate, temperature, and oxygen saturation.
13. Prepare to assess lab values as ordered.
14. Ensure oxygen and suction equipment are present and functioning.
15. Implement measures to decrease stress level, such as provision of a quiet environment and low lighting.
16. Monitor temperature per department protocol.
17. Assess intake and output (I&O) every 1 hour.

ANTEPARTUM ONGOING ASSESSMENT:

Goals of patient management are:

1. Early recognition of severe or worsening preeclampsia or development of eclampsia.
2. Prolongation of pregnancy to optimize fetal maturation must be weighed against risks of pregnancy continuation.

Mild Preeclampsia:

1. Obtain blood pressure, pulse, respirations, and oxygen saturation every 4 hours.
2. Assess lung sounds every 4 hours.
3. Assess deep tendon reflexes (DTR's), Clonus, edema, level of consciousness (LOC), headache (HA) visual disturbances, epigastric pain every 8 hours.
4. Obtain Non Stress Test (NST) or monitor Fetal Heart Rate (FHR) with uterine activity for 30 minutes every shift or as condition warrants.
5. Assess fetal movement every shift.

Severe Preeclampsia:

1. Obtain blood pressure, pulse, respirations, and oxygen saturation hourly.

2. Assess lung sounds every 2 hours.
3. Assess deep tendon reflexes (DTR's), Clonus, edema, level of consciousness (LOC), Headache (HA) visual disturbances, epigastric pain every 4 hours.
4. Monitor FHR and uterine activity continuously.

INTRAPARTUM ONGOING ASSESSMENT:

Mild Preeclampsia:

1. Obtain blood pressure, pulse, respirations, and oxygen saturation every 60 minutes.
2. Assess lung sounds every 4 hours.
3. Assess deep tendon reflexes (DTR's), clonus, edema, level of consciousness (LOC), headache (HA) visual disturbances, epigastric pain every 8 hours.
4. Monitor FHR and uterine activity continuously.

Severe Preeclampsia:

1. Obtain blood pressure, pulse, respirations, and oxygen saturation every 30 minutes.
2. Assess lung sounds every 2 hours.
3. Assess Deep Tendon Reflexes (DTR's), clonus, edema, level of consciousness (LOC), headache (HA) visual disturbances, epigastric pain every 4 hours.
4. Monitor FHR and uterine activity continuously.

POSTPARTUM TO DISCHARGE ONGOING ASSESSMENT:

Mild Preeclampsia:

1. Obtain blood pressure, pulse, respirations, and oxygen saturation every 4 hours.
2. Assess lung sounds every 4 hours.
3. Assess deep tendon reflexes (DTR's), Clonus, edema, level of consciousness (LOC), headache (HA) visual disturbances, epigastric pain every 8 hours.

Severe Preeclampsia:

1. Obtain blood pressure, pulse, respirations, and oxygen saturation every 60 minutes for first 24 hours after delivery then every 4 hours.
2. Assess lung sounds every 2 hours for first 24 hours after delivery then every 4 hours.
3. Assess deep tendon reflexes (DTR's), clonus, edema, level of consciousness (LOC), headache (HA) visual disturbances, epigastric pain every 4 hours.

MAGNESIUM SULFATE:

Magnesium sulfate is administered as a first line drug to prevent maternal eclamptic seizures. (See Magnesium Sulfate section, pg. 48)

ANTIHYPERTENSIVES:

Background:

1. A sustained diastolic pressure greater than 160 mmHg systolic OR greater than 105 mmHg diastolic is treated with IV antihypertensive medication to protect the patient from cerebral vascular accident.
2. The goal is a diastolic pressure of 90-100mm Hg to maintain perfusion.
3. Labetalol is a combined alpha and beta-blocker, resulting in decreased peripheral vascular resistance without altering heart rate or cardiac output. Its use is contraindicated in patients with bronchial asthma, heart block and severe bradycardia.
4. Hydralazine is a vasodilator and results in vasodilation of vascular smooth muscle.

Administration:

1. Ensure presence of mainline IV infusion.
2. Monitor the fetal heart rate continuously if a viable fetus is present.
3. Maintain bedrest during and for 3 hours following medication administration. Assess for postural hypotension prior to ambulation.
4. If unable to control blood pressure, contact physician re: consideration of other medications and/or transfer to a higher level of care.
5. Hydralazine (Apresoline):
 - a. Administer initial dose IV push over 1-2 minutes. (usual dose range is 5-10 mg)
 - b. May repeat dose at 20-minute intervals until desired blood pressure is achieved or a cumulative dose of 30-40 mg is reached
6. Labetalol:
 - a. IV Push:
 - i. Administer initial dose IV push over 2 minutes. (usual dose is 10-20 mg)
 - ii. Repeat doses may be given at 10 minute intervals.
 - b. Continuous IV:
 - i. Consider collaborative care with intensive care unit.
 - ii. Initiation of continuous cardiac monitoring.
 - iii. Infuse a continuous labetalol infusion pump until diastolic pressure is 90-100 mmHg.
 - c. Maximum dose is 300 mg/24 hours.

Reportable Conditions:

1. Notify provider for:
 - a. Diastolic blood pressure less than 80 or greater than 105 following medication administration.
 - b. Category II or III fetal heart rate tracing following antihypertensive administration.
 - c. Sustained maternal heart rate less than 50 bpm or greater than 120 bpm during or within 30 minutes following medication administration.

ECLAMPSIA MANAGEMENT:

Background:

- Eclampsia is characterized by convulsions and loss of consciousness, which can occur without warning during the antepartum, intrapartum or postpartum period.
- The eclamptic patient is at risk for aspiration and cerebral hemorrhage.
- Fetal bradycardia frequently occurs during and following an eclamptic seizure.
- Best treatment for baby is maternal stabilization.

MANAGEMENT:

1. Notify charge nurse, attending provider, and anesthesiologist/CRNA immediately. Initiate emergency pager (if institution has instituted).
2. Position patient on side.
3. Protect from injury.
4. Prepare to administer magnesium sulfate.
5. Anticipate obtaining lab tests (magnesium level, blood for liver enzymes, kidney function, etc.).
6. Following seizure:
 - a. Suction mouth.
 - b. Give oxygen by non-rebreather mask at 10 liters per minute.
 - c. Provide ventilatory support as needed.
 - d. Assess blood pressure, pulse, and respirations every 5 minutes.
 - e. Assess oxygen saturation and level of consciousness every 15 minutes until stable for a minimum of one hour.
 - f. Monitor fetal heart rate and uterine activity continuously if viable fetus is present.
 - g. Observe for signs and symptoms of placental abruption or impending delivery.
 - h. Obtain order for indwelling catheter.

Appendix V: Daily Assessment for Delivery versus Continuing Pregnancy

Criteria:	Present	
	Yes	No
Persistent maternal headache	Yes	No
Visual disturbance (blurred or scotomata)	Yes	No
Hypoxia (O2 saturation <95%) or pulmonary edema on clinical exam	Yes	No
Persistent BP >160mmHg systolic or >105 mmHg despite medical management	Yes	No
Oliguria (<500 ml/24 hours)	Yes	No
Evidence of renal failure (serum Creatinine >1.2 mg/dL)	Yes	No
Thrombocytopenia (platelet count <100,000/mm ³)	Yes	No
Elevated ALT>70 U/L	Yes	No
Evidence of hemolysis (LDH>600, bilirubin>1.2 mg/dL or abnormal peripheral blood smear)	Yes	No
Abnormal coagulation (elevated PT/PTT or fibrinogen <300)	Yes	No
Abnormal Fetal NST and/or BPP	Yes	No
	Yes To ANY of above CONSIDER DELIVERY	No To ALL of above CONTINUE PREGNANCY

Appendix W: Classification of Evidence Grading

Evidence Categories

Type of Study or Evidence	
I	Evidence obtained from at least one properly designed randomized controlled trial.
II-1	Evidence obtained from well-designed controlled trials without randomization.
II-2	Evidence obtained from well-designed cohort or case-control analytic studies, preferably from more than one center or research group.
II-3	Evidence obtained from multiple time series with or without intervention. Well done QI studies with statistical process control analyses or the like fall into this category. Dramatic results in uncontrolled experiments also could be regarded as this type of evidence.
III	Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

Level of Recommendations	
A	Recommendations based on high quality and consistent evidence.
B	Recommendations based on limited or inconsistent evidence.
C	Recommendations based primarily on consensus and expert opinion.

Adapted from USPSTF and ACOG